EPBC Referral Supplementary Report No. 5

Boskalis Cambridge Gulf Marine Sand Proposal Western Australia

RESPONSE TO REQUEST FOR FURTHER INFORMATION (RFI)









Prepared for Boskalis Australia Pty Ltd by EcoStrategic Consultants

In support of Project Referral under Part 7 of Commonwealth *Environment Protection* & *Biodiversity Conservation Act*

AUGUST 2025





CONTENTS

PROPONENT CONTACTS	2
REFERRAL DOCUMENTATION	3
ACRONYMS	5
PROJECT LOCATION	6
1. BACKGROUND & PURPOSE OF THIS REPORT	7
2. SUMMARY RESPONSES TO RFI	7
3. RESPONSE TO RFI ITEM 1 - LISTED THREATENED SPECIES	11
3.1 Potential for Significant Impacts	11
3.2 Diseases & Pathogens	12
3.3 Species Conservation Advice	14
3.4 Species Recovery Plans	36
3.5 Species Threat Abatement Plans	48
4. RESPONSE TO RFI ITEM 2 - ECONOMIC & SOCIAL MATTERS	48
5. RESPONSE TO RFI ITEM 3 - ENVIRONMENTAL MANAGEMENT PLAN	48
REFERENCES	49
ANNEX 1: DCCEEW RFI	50

PROPONENT CONTACTS

Alwin van den Bosch General Manager

Boskalis Australia Pty Ltd

Suited 1.3 / 9 Havelock St, West Perth 6005 Tel 08 9327 1000 alwin.vd.bosch@boskalis.com

Steve Raaymakers

Consultant to Boskalis

EcoStrategic Consultants

PO Box 968, Edge Hill, Cairns 4870

Mobile 040 9909 422

steve@eco-strategic.com

REFERRAL DOCUMENTATION

This report is part of a larger set of documents submitted as part of Boskalis Australia's referral under Part 7 of the Commonwealth *Environment Protection & Biodiversity Conservation Act* (EPBC Act), as listed in the table below.

Doc	Reference	Electronic File Names (PDFs) (except Doc No.s 9 & 10 which are Excel files).
No.	(Author/yr)	As required, these file names are how the reports are referenced in the online referral submitted via the EPBC Act Business Portal https://epbcbusinessportal.environment.gov.au
0	-	EPBC Referral - Boskalis Cambridge Gulf - List of Preliminary Documents.
1	BKA (2024a)	EPBC Referral Report No. 1 - Boskalis Cambridge Gulf - Description of Proposed Action & Regulatory Framework.
2	BKA (2024b)	 EPBC Referral Report No. 2 - Boskalis Cambridge Gulf - Setting & Existing Environment. Includes in same document: Annex 3 - Drop Camera Video Extracts. Annex 4 - Dry Season Sample Point Specs. Annex 5 - Wet Season Sample Point Specs. Annex 6 - Benthic Taxa per Sample Point - Dry Season Maps. Annex 7 - Benthic Taxa per Sample Point - Wet Season Maps. Annex 8 - Benthic Taxa per Sample Point - Dry Season Graphs. Annex 9 - Benthic Taxa per Sample Point - Wet Season Graphs. Annex 11 - Sediment Contamination Assessment. Annexes 1, 2, 10, 12, 13 and 14 are submitted as separate documents as listed below.
3	BKA (2024c)	EPBC Referral Report No. 2 - Boskalis Cambridge Gulf - Annex 1 - Sand Assessment.
4	MScience (2024)	EPBC Referral Report No. 2 - Boskalis Cambridge Gulf - Annex 2 - MScience BCH Methods.
5	Sensorem (2024)	EPBC Referral Report No. 2 - Boskalis Cambridge Gulf - Annex 10 - Aerial Drone Lidar Report.
6	Price & Raaymakers (2024)	EPBC Referral Report No. 2 - Boskalis Cambridge Gulf - Annex 12 - Cape Domett Turtle Data Report.
7	Univ. Canberra (2024)	EPBC Referral Report No. 2 - Boskalis Cambridge Gulf - Annex 13 - Marine eDNA Report.
8	BKA (2024d)	 EPBC Referral Report No. 2 - Boskalis Cambridge Gulf - Annex 14 - Marine Mega-fauna Surveys Report. Includes in same document: Appendix 1 - MMF Sightings Master Data Tables. Appendix 2 - MMF Images. Appendix 3 - MMF Sighting Locations. Appendices 4 and 5 are submitted as separate Excel files as listed below.
9	BKA (2024e)	EPBC Referral Report No. 2 - Boskalis Cambridge Gulf - Annex 14 - Appendix 4 - Species Data - Dry Season (Excel).
10	BKA (2024f)	EPBC Referral Report No. 2 - Boskalis Cambridge Gulf - Annex 14 - Appendix 5 - Species Data - Wet Season (Excel).
11	BKA (2024g)	EPBC Referral Report No. 3 - Boskalis Cambridge Gulf - Traditional Owner Matters. Includes in same document: • Annex 1 - BAC Native Title Determination Map. • Annex 2 - MG Native Title Determination Map. • Annex 3 - Letter from BAC.

Dos	Doforon	Floatronia Fila Names (PDFs) (except Dec No. c. 0. 9. 40 which are Excel Files)
Doc No.	Reference (Author/yr)	Electronic File Names (PDFs) (except Doc No.s 9 & 10 which are Excel files). As required, these file names are how the reports are referenced in the online referral submitted via the EPBC Act
		Business Portal https://epbcbusinessportal.environment.gov.au
		Annex 4 - Letter from MG.
12	ВКА	EPBC Referral Report No. 4 - Boskalis Cambridge Gulf - Impact Assessments. Includes in same document:
	(2024h)	Annex 1 - Main Datasets Used to Inform Impact Assessments.
		Annex 2 - Shipping & Oil Spill Risk Assessment. Annex 3 - Plume Mitigation Capability Statement.
		Annex 4 - Marine Mega-fauna Capability Statement.
13	PCS	EPBC Referral Report No. 5 - Boskalis Cambridge Gulf - Metocean & Sed Dynamics Initial Report.
	(2024a)	Includes in same document <u>Annex 1</u> - Supplementary Technical Note.
		Annex 2 is submitted as a separate document as listed below.
14	PCS	EPBC <u>Referral Report No. 5</u> - Boskalis Cambridge Gulf - <u>Annex 2</u> - <i>Factual Data Report</i> .
	(2024b)	(NOTE: Superseded by Updated Factual Data Report - see Doc No. 19, Referral Report No. 8 - Annex B below).
15	BKA (2024i)	EPBC <u>Referral Report No. 6</u> - Boskalis Cambridge Gulf - <i>Consultation</i> .
		Includes in same document Annex 1 - List of Meeting Minutes.
16	BKA (2024j)	EPBC Referral Report No. 7 - Boskalis Cambridge Gulf - Commonwealth Matters.
	(,j,	Includes in same document Annex 1 - PMST Report for POA & 10 Km Buffer.
17	PCS	EPBC Referral Report No. 8 - Boskalis Cambridge Gulf - Metocean & Sed Dynamics Full Modelling Report.
	(2025a)	Appendices and Annexes are submitted as a separate document each, as listed below.
18	PCS	EPBC Referral Report No. 8 - Boskalis Cambridge Gulf - Appendices.
	(2025b)	Appendix A - Model Calibration and Validation Plots.
		Appendix B - Hydrodynamic and Wave Impact Plots. Appendix C - Sediment Transport Impact Plots.
		Appendix D - Sediment Plume Modelling Results.
19	PCS	EPBC Referral Report No. 8 - Boskalis Cambridge Gulf - Annexes.
	(2025c)	Annex A - Independent Expert Review.
		Annex B - Updated Factual Data Report.
NOTE	: The documen	its listed above were submitted in the initial referral. The documents listed below were submitted after the initial referral.
20	Nocterra	EPBC Referral Supplementary Report No. 1 - Boskalis Cambridge Gulf - Light Assessment.
	(2025)	
21	Resonate	EPBC Referral Supplementary Report No. 2 - Boskalis Cambridge Gulf - Noise Assessment.
	(2025)	
22	ВКА	EPBC Referral Supplementary Report No. 3 - Boskalis Cambridge Gulf - Commonwealth Environmental Management
	(2025a)	Plan (C-EMP).
23	ВКА	EPBC Referral Supplementary Report No. 4 - Boskalis Cambridge Gulf - Additional Information.
	(2025b)	Current Speeds in the POA & Turtle Swimming Speeds.
		Analysis of Turtle Satellite Tracking - Cape Domett. Replacin Complitity Shoot Trailer System Hopers Produces
		Boskalis Capability Sheet - Trailer Suction Hopper Dredgers.
24	ВКА	THIS DOCUMENT: EPBC Referral Supplementary Report No. 5 - Boskalis Cambridge Gulf - Response to Request
	(2025c)	for Further Information.

ACRONYMS

BIA Biologically Important Area
BKA Boskalis Australia Pty Ltd

BWM Convention International Convention for the Control & Management of Ships' Ballast Water & Sediments

C-EMP Commonwealth Environmental Management Plan (as presented in Supplementary Report No. 3)

CEO Commonwealth Environmental Outcome (as contained in the C-EMP)

CG Cambridge Gulf

CMS Convention on Migratory Species

DAFF Commonwealth Department of Agriculture, Fisheries & Forestry
DBCA WA Department of Biodiversity, Conservation & Attractions

DCCEEW Commonwealth Department of Climate Change, Energy, the Environment & Water

DEMIRS WA Department of Energy, Mines, Industry Regulation & Safety

EIS Environmental Impact Statement (under EPBC Act)

EPA WA Environmental Protection Authority
EP Act WA Environmental Protection Act

EPBC Act Commonwealth Environment Protection & Biodiversity Conservation Act

IMO International Maritime Organization

LiDAR Light Detection & Ranging

MARPOL International Convention for the Prevention of Pollution from Ships

MFOA Marine Fauna Observation and Avoidance

MNES Matters of National Environmental Significance (under Commonwealth EPBC Act)

PMST (Commonwealth) Protected Matters Search Tool
Ramsar Convention on Wetlands of International Importance

RFI Request for Further Information

SPV Sand Production Vessel
TO Traditional Owner

TSHD Trailer Suction Hopper Dredger WA Western Australia (State of)

PROJECT LOCATION




FIGURE 1: Location of the proposed action in Cambridge Gulf near Wyndham in the northeast of WA.

1. BACKGROUND & PURPOSE OF THIS REPORT

- Boskalis Australia Pty Ltd (BKA) is assessing the feasibility of developing a marine sand-sourcing operation (the proposed action) in Cambridge Gulf (CG) near Wyndham in the northeast of Western Australia (WA) (Figure 1). BKA currently holds two sand exploration tenements in CG under the WA *Mining Act*, as the basis for the proposed action.
- 2. A detailed description of the proposed action is presented in <u>EPBC Referral Report No. 1 Description of the Proposed Action & Regulatory Framework</u> and is not repeated in this report for reasons of economy.
- 3. To support its assessment BKA has undertaken a wide range of comprehensive studies since 2018. These studies find that the proposed action is feasible and viable and unlikely to cause significant environmental impacts, as defined under the WA Environmental Protection Act (EP Act) and the Commonwealth Environmental Protection & Biodiversity Conservation Act (EPBC Act). Never-the-less, as a responsible company with stringent environmental and social policies, BKA self-referred the proposal to both the State and the Commonwealth under their respective Acts, for their determination of what further environmental assessments might be required, if any. The EPBC Act referral was submitted in January 2025.
- 4. Subject to the outcomes of the State and Commonwealth EPBC Act referral processes, BKA plans to apply to the WA Department of Energy, Mines, Industry Regulation & Safety (DEMIRS) to convert a reduced part of the two Exploration Tenements to a single Mining Tenement, shown as the 'proposed operational area' (POA) on Figure 1.
- 5. On 27 June 2025 a delegate of the Commonwealth Minister for the Environment decided that:
 - a) the proposed action is a controlled action under the EPBC Act,
 - b) it will be assessed by preliminary documentation; and
 - c) further information was required to assess relevant impacts of the proposed action.
- 6. On 16 July 2025 DCCEEW issued a letter to BKA with a Request for Further Information (RFI), under section 95A(2) of the EPBC Act, as attached in Annex 1.
- 7. The purpose of this report is to provide BKA's response to the RFI.

2. SUMMARY RESPONSES TO RFI

- 1. Annex 1 contains DCCEEW's letter of 16 July 2025 with the RFI requirements. Table 1 presents each item and sub-item of the RFI along with a summary of BKA's response to each, with reference to the more detailed responses in the following sections below. The item numbering has been introduced by BKA to assist in organizing the responses, and are not used in DCCEEW's RFI letter. However, the headings and contents are exactly the same.
- 2. There is considerable repetition of material in the responses to each item of the RFI. This is because responses to all items have been included and addressed in turn, so as to provide a complete picture of how BKA has addressed all aspects of the RFI. Where possible, repetition has been reduced by referencing other sections of this report and relevant sections of the various Referral Reports already submitted, where the relevant responses have already been provided.

TABLE 1: Summary responses to RFI

DCCEEW Request	Summary BKA Response
Item 1: Listed threatened species (sections 18 & s 18A of EPBC Act)	
Item 1.1: Potential significant impacts:	Refer section 3.1 below for detailed response.
The department considers that the proposed action is likely to have a significant impact on: - Flatback Turtle (Natator depressus) – Vulnerable (Migratory). - Australian Snubfin Dolphin (Orcaella heinsohni) – Vulnerable (Migratory). - Australian Humpback Dolphin (Sousa sahulensis) – Vulnerable (Migratory).	BKA does not understand on what bases the department considers that the proposed action is <i>likely to have a significant impact</i> on the three listed species, as described by the EPBC Act significant impact criteria, and considering the nature of the proposed operation and proposed impact prevention, mitigation, monitoring and response measures. BKA has sought clarification on this from the department, which has not been received. BKA's systematic and scientific impact assessment, conducted in accordance with the EPBC Act significant impact criteria and the impact mitigation hierarchy, as presented in BKA's referral reports, finds that it is highly unlikely that the proposed action would cause significant or residual or irreversible impacts on the listed species. This is further supported by the information submitted in this response to the RFI.
Item 1.2: Diseases & pathogens:	Refer section 3.2 below for detailed response.
The department notes that i) the Marine bioregional plan for the North-west Marine Region, ii) the Conservation Advice for Orcaella heinsohni (Australian snubfin dolphin), iii) the Conservation Advice for Sousa sahulensis (Australian humpback dolphin) and iv) the Recovery Plan for Marine Turtles in Australia identify disease (and pathogens) as pressures/threats to these species. Please provide further discussion of this threat, together with management measures (mittigation, early-warning monitoring, research programs) aimed at early detection of new diseases affecting populations of the threatened species mentioned above. Please discuss these threats in the Preliminary Documentation and in the EMP. Note: This is separate and additional to the measures proposed in EO 4 - "no marine pest species are introduced via the SPV's ballast water discharges or hull bio-fouling".	The Marine bioregional plan for the North-west Marine Region is not relevant as it applies to Commonwealth waters only, while CG is within internal State waters, landward of the baseline and thus not even in 3nm coastal State waters. The Marine bioregional plan contains a single reference to 'disease' as being a potential 'pressure' on biodiversity of the region, with potential sources of disease being identified as aquaculture, fishing, tourism and shipping (with only the latter being relevant to the project, as the SPV is a ship). Potential diseases from shipping relate to ballast water discharges, which have been fully addressed in accordance with both IMO and Commonwealth requirements. BKA is not aware of any other potential vectors / mechanisms whereby the SPV could cause introductions of diseases of the listed species. BKA has sought advice from DCCEEW on examples of other similar marine projects in Australia where diseases and pathogens have been an issue, and the management measures that have been required by DCCEEW in order to address this - so that BKA can follow established best-practices. Such advice has not been received. The Recovery Plan for Marine Turtles in Australia contains a dedicated section on diseases and pathogens, and implies that diseases in marine turtles are natural but may be exacerbated by poor water quality. The project will not have negative impacts on water quality. The Recovery Plan also states that 'To date, there are no recorded occurrences of diseases and pathogens affecting the viability of a marine turtle stock in Australia.' DCCEEW's Conservation Advice for Snubfin Dolphins and for Humpback Dolphins similarly imply potential links between poor water quality and dolphin skin diseases. As above, the project will not have negative impacts on water quality. Some of the documents do make specific reference to bacterial infections which can result from injuries caused by vessel strikes – this is addressed through measures to prevent and mitigate vessel strikes.

DCCEEW Request	Summary BKA Response
Item 1.3: Consideration of relevant conservation advice, recovery plans and/or threat abatement plans: Please demonstrate that the proposal has had regard to relevant conservation advices, and is not inconsistent with recovery plans and/or threat abatement plans, including but not limited to those listed in Annex 1 of Attachment B.	Much of this was addressed in BKA's Referral Reports as submitted, and especially: - EPBC Referral Report No. 2 - Setting & Existing Environment Section 9 - Marine Fauna - EPBC Referral Report No. 2 - Setting & Existing Environment - ANNEX 14 - Marine Fauna Surveys Report. - EPBC Referral Report No. 4 - Impact Assessments Section 10 - Impact Assessment - Marine Fauna, - EPBC Referral Report No. 7 - Commonwealth Matters Section 10 - Potential Impacts on Species-based MNES. - EPBC Supplementary Report No. 1 - Light Assessment EPBC Supplementary Report No. 2 - Noise Assessment. Section 3.3. below assesses the following seven Conservation Advice documents: - Conservation Advice for Orcaella heinsohni (Australian Snubfin Dolphin), March 2025 Conservation Advice for Sousa sahulensis (Australian Humpback Dolphin), March 2025 Conservation Advice for Glyphis glyphis (Speartooth Shark), April 2014 Conservation Advice for Glyphis garricki (Northern River Shark). April 2014 Conservation Advice for Pristis pristis (Largetooth Sawfish), April 2014 Conservation Advice for Pristis zijsron (Green Sawfish), undated Conservation Advice for Pristis zijsron (Green Sawfish), October 2009. Section 3.4. below assesses the following two Species Recovery Plans: - Sawfish & River Sharks Multi-species Recovery Plan, 2015 Recovery Plan for Marine Turtles in Australia, 2017-2027.
	Section 3.5 below assesses the following Threat Abatement Plan: — Threat Abatement Plan for the impacts of marine debris on the vertebrate wildlife of Australia's coasts and oceans (2018) (short title Marine Debris Threat Abatement Plan).
Item 2: Economic and social matters	Refer section 4 below.
Please provide further detail on the social and economic costs and/or benefits of undertaking the proposed action, including:	DCCEEW has since advised that the information on economic and social matters contained in the Referral Reports as already submitted is adequate and no further response is required on this item.
 An estimate of any anticipated economic costs and/or benefits (in AUD), particularly with reference to the domestic market. 	
The basis for any estimations of costs and/or benefits.	
 Detail any social matters around the proposed action. This includes benefits to the local Traditional Owners. 	
 Any potential employment opportunities expected to be generated by the proposed action, including any potential opportunities for local Traditional Owners groups. 	
Any funding the project has received from federal departments or agencies.	

DCCEEW Request	Summary BKA Response
Item 3: Environmental Management Plan (EMP)	Refer section 5 below.
A separate Attachment B was provided by DCCEEW with detailed comments on the Draft EMP that had been submitted to them several weeks previously.	Draft 2 of the proposed EMP has been developed, addressing DCCEEW's review comments on Draft 1.
Responses to each comment are provided in Attachment B - DCCEEW Comments on the EMP v.1 - BK Responses.	This is submitted separately as <u>EPBC Supplementary Report No. 3 – Commonwealth Environmental Management Plan (C-EMP)</u> , along with a table listing how each of DCCEEW's review comments have been addressed.

3. RESPONSE TO RFI ITEM 1 - LISTED THREATENED SPECIES

3.1 Potential for Significant Impacts

- 1. BKA does not understand on what bases DCCEEW considers that the proposed action is likely to have a significant impact on the three listed species (Flatback Turtle, Snubfin Dolphin and Humpback Dolphin), as described by the <u>EPBC Act significant impact guidelines</u> (DCCEEW 2021), and considering the nature of the proposed operation and associated impact-causing mechanisms, and proposed impact prevention, mitigation, monitoring and response measures. BKA has sought clarification on this from the department, which has not been received.
- 2. BKA's systematic and scientific impact assessments, conducted in accordance with the EPBC Act significant impact criteria and the impact mitigation hierarchy, as presented in BKA's Referral Reports, finds that it is highly unlikely that the proposed action would cause significant, residual or irreversible impacts on the listed species. This is further supported by the information submitted in this response to the RFI.
- 3. BKA's systematic impact assessments with regard to the three listed species are detailed in the following Referral Reports, and the details are not repeated in this report, for reasons of economy.

a) Flatback Turtles:

- EPBC Referral Report No. 2 Setting & Existing Environment.
 - Section 9.4 presents a detailed description of marine turtles, including Flatback Turtles in the CG area, supported by Annex 12 to that report - <u>DBCA Cape Domett Turtle Data Report</u>.
- EPBC Referral Report No. 4 Impact Assessments.
 - Potential impacts of the proposed operation on Flatback Turtles are assessed in Section 10.3.2 of in accordance with WA EPA guidelines and impact mitigation hierarchy, and find no significant or residual impacts in accordance with those guidelines.
- EPBC Referral Report No. 7 Commonwealth Matters.
 - Potential impacts of the proposed operation on Flatback Turtles are assessed in Section 10.2 of in accordance with EPBC Act significant impact criteria and impact mitigation hierarchy, and finds no significant or residual impacts in accordance with those criteria.
- EPBC Supplementary Report No. 1 Light Assessment
 - Potential impacts of light emissions from the SPV on nesting and hatching turtles in the CG area are assessed in accordance with both the National Light Pollution Guidelines for Wildlife (DCCEEW 2023) and WA EPA requirements, and finds no significant impacts.
- EPBC Supplementary Report No. 2 Noise Assessment
 - This includes detailed modelling of predicated noise emissions from the SPV and assessment of potential auditory injury and behavioural impacts on marine turtles, in accordance with the US NMFWS criteria (as required by WA EPA), using a risk assessment approach, and finds that potential impacts are negligible.
- EPBC Supplementary Report No. 3 Commonwealth Environmental Management Plan (C-EMP)
 - This includes best practice impact avoidance, mitigation, monitoring and response actions for marine turtles in accordance with the impact mitigation hierarchy.
- EPBC Referral Supplementary Report No. 4 Additional Information.
 - Additional information on marine turtle issues is presented in Section 2 Current Speeds in the POA & Turtle Swimming Speeds, and Section 3 - Analysis of Turtle Satellite Tracking - Cape Domett.

Further assessment is provided for marine turtles in Table 10 in section 3.4 below, which assesses how BKA has taken account of the Recovery Plan for Marine Turtles.

- b) <u>Snubfin & Humpback Dolphins</u> (combined as the assessments are the same):
 - EPBC Referral Report No. 2 Setting & Existing Environment.
 - Section 9 Marine Fauna describes marine fauna in the area based on all available information and the site surveys commissioned by BKA.

 Section 9.4.1 - Australian Snubfin Dolphin and Section 9.4.2 - Australian Humpback Dolphin provide specific descriptions of these species in the area, based on all available information and the site surveys commissioned by BKA.

- EPBC Referral Report No. 2 - Setting & Existing Environment - ANNEX 14 - Marine Fauna Surveys Report.

This presents the methods and results of marine fauna surveys commissioned by BKA, including for Snubfin and Humpback Dolphins, including literature review, consultations with relevant experts and stakeholders, assessment of previous surveys conducted by others in the area (e.g Brown et al 2016 & 2017), and the dry- and wet-season surveys carried out in accordance with the National Guidelines for the Survey of Cetaceans, Marine Turtles and Dugong (DCCEEW 2024).

- EPBC Referral Report No. 4 - Impact Assessments.

- Section 10 Impact Assessment Marine Fauna, assesses potential impacts of the proposal on marine fauna in accordance with WA EPA guidelines and impact mitigation hierarchy.
- Section 10.3.1 Snubfin & Humpback Dolphins applies the impact assessment to these species and finds no significant or residual impacts in accordance with WA EPA guidelines and impact mitigation hierarchy.

- EPBC Referral Report No. 7 - Commonwealth Matters.

 Section 10.3 - Specific Assessment for Snubfin Dolphins assesses potential impacts of the proposal on this species in accordance with the EPBC Act significant impact criteria and impact mitigation hierarchy, and finds no significant or residual impacts in accordance with these criteria. It applies equally to Humpback Dolphins given the similarity of these two species.

- EPBC Supplementary Report No. 2 - Noise Assessment.

 This includes detailed modelling of predicated noise emissions from the SPV and assessment of potential auditory injury and behavioural impacts on Snubfin and Humpback Dolphins, in accordance with the US NMFWS criteria (as required by WA EPA), using a risk assessment approach, and finds that potential impacts are negligible.

- EPBC Supplementary Report No. 3 - Commonwealth Environmental Management Plan (C-EMP)

 This includes best practice impact avoidance, mitigation, monitoring and response actions for Snubfin and Humpback Dolphins in accordance with the impact mitigation hierarchy.

Further assessment is provided for Snubfin Dolphins in Table 3 and for Humpback Dolphins in Table 4 in section 3.3 below, which assess how BKA has taken account of the Conservation Advice documents for these species.

4. When considering potential impacts under the EPBC Act, it is important to give due consideration to the listed threatened species criteria under the EPBC Act significant impact guidelines (DCCEEW 2021). As can be seen from EPBC Referral Report No. 7 - Commonwealth Matters, the significant impact guidelines have a focus on conservation at the population level, and it is clear from the assessments in that report that the proposed action is not likely to present a risk of impacts at the population level. Any suggestion to the contrary should be supported with scientific explanation of the bases for the assessment, considering the nature of the proposed operation and associated impact-causing mechanisms, and the proposed impact prevention, mitigation, monitoring and response measures.

3.2 Diseases & Pathogens

- 1. Like all living biota (plants and animals), marine fauna including dolphins and turtles can suffer from a wide range of diseases and pathogens. As for all living biota, diseases are generally the result of natural causes. However, in some circumstances they can be exacerbated by human causes, for example a reduction in water quality from pollution from land-based industry, which can lower the resistance and immunity of marine plants and animals to disease, or physical injury to animals that create wounds followed by bacterial infection (usually non-fatal). Marine diseases and pathogens can also be translocated by humans and introduced to new areas, for example via the transfer of aquaculture stock that might carry diseases and pathogens to new areas, or via ballast water discharges from vessels.
- 2. As outlined in section 2 above, item 1.2 of DCCEEW's RFI states:

'The department notes that:

- i) the Marine bioregional plan for the North-west Marine Region,
- ii) the Conservation Advice for Orcaella heinsohni (Australian snubfin dolphin),
- iii) the Conservation Advice for Sousa sahulensis (Australian humpback dolphin); and
- iv) the Recovery Plan for Marine Turtles in Australia;

identify disease (and pathogens) as pressures/threats to these species.

Please provide further discussion of this threat, together with management measures (mitigation, early-warning monitoring, research programs) aimed at early detection of new diseases affecting populations of the threatened species mentioned above.'

- 3. The Marine bioregional plan for the North-west Marine Region is not relevant to the proposed action as it applies to Commonwealth waters only, while CG is within internal State waters, landward of the baseline and thus not even in 3 nm coastal State waters.
- 4. The *Marine bioregional plan* contains a single reference to 'disease' as being a potential 'pressure' on biodiversity of the region, with potential sources of disease being identified as aquaculture, fishing, tourism and shipping (with only the latter being relevant to the project, as the SPV is a ship).
- 5. Potential diseases from shipping relate to ballast water discharges, which have been fully addressed in the Referral Reports and the proposed EMP in accordance with both IMO and Commonwealth requirements, as per Commonwealth Environmental Outcome (CEO) 6 Marie Pests in EPBC Supplementary Report No. 3 Commonwealth Environmental Management Plan (C-EMP).
- 6. As far as BKA can determine, there are no other potential vectors / mechanisms whereby the SPV could cause introductions of diseases of the listed species into CG. The seabed in the POA where sand-sourcing is proposed, comprises highly dynamic sand-waves, which are constantly mobilised, suspended and mixed by the extreme tidal currents in CG, so nothing could be buried in them, including disease and pathogens. A detailed description of the environmental conditions in the POA is provided in sections 5, 6.4.2, 6.4.3, 6.4.4 and 6.4.4.8 of EPBC Referral Report No. 2 Setting & Existing Environment.
- 7. As described in Annex 11 Sediment Contamination Assessment of <u>EPBC Referral Report No. 2 Setting & Existing Environment</u>, the seabed sediments in the POA are free of contaminants as assessed in accordance with the National Assessment Guidelines for Dredging (NAGD) 2009.
- 8. BKA has sought advice from DCCEEW on examples of other similar marine projects in Australia where diseases and pathogens have been an issue for dolphin and marine turtles, and the management measures that have been required by DCCEEW in order to address this so that BKA can follow established best-practices. Such advice has not been received.
- 9. The Recovery Plan for Marine Turtles in Australia contains a dedicated section on diseases and pathogens, and implies that natural diseases in marine turtles may be exacerbated by poor water quality, primarily from pollution from land-based industry and other sources. The project will not have negative impacts on water quality, as assessed in accordance with WA EPA guidelines on marine environmental quality and the impact mitigation hierarchy, in Section 9 Impact Assessment Marine Environmental Quality of EPBC Referral Report No. 4 Impact Assessments.
- 10. The Recovery Plan also states that disease outbreaks in food sources, such as seagrass, can indirectly affect the health of marine turtles. This is not a risk for this proposed action as there are no seagrasses in CG and the area does not provide foraging habitat for marine turtles, as described in Section 6 Benthic Communities & Habitats, of EPBC Referral Report No. 2 Setting & Existing Environment.
- 11. The Recovery Plan also states that '<u>To date, there are no recorded occurrences of diseases and pathogens affecting the viability of a marine turtle stock in Australia</u>.' As far as BKA can determine, there are no operational mechanisms whereby the operation of the SPV in CG would change this situation, by affecting the viability of the marine turtle stock in the area through diseases and pathogens.
- 12. DCCEEW's Conservation Advice for Snubfin Dolphins and for Humpback Dolphins similarly states potential links between poor water quality and exacerbation of natural skin diseases in dolphins. As above, the proposed operation will not have negative impacts on water quality, so this is not a risk for this project.
- 13. Some of the documents make specific reference to bacterial infections which can result from injuries caused by vessel strikes this is addressed through measures to prevent and mitigate vessel strikes, as per CEO 7 Vessel Strikes in EPBC Supplementary Report No. 3 Commonwealth Environmental Management Plan (C-EMP).
- 14. Overall, the only aspects of the operation which could have relevance to diseases and pathogens in dolphins and marine turtles in CG are ballast water discharges and potential vessel strikes causing injury, and thus potential infection, in an individual animal. Comprehensive impact prevention, mitigation, monitoring and response actions are proposed for both of these aspects, as outlined above.
- 15. Early-warning monitoring and research programs aimed at early detection of 'new' diseases is a responsibility for conservation management agencies, veterinary authorities and research bodies, and not for a commercial operation that does not present the risk of introducing new diseases.

3.3 Species Conservation Advice

- 1. As outlined in section 2 above, item 1.3 of DCCEEW's RFI states 'Please demonstrate that the proposal has <u>had regard to relevant conservation advices</u>, and is not inconsistent with recovery plans and/or threat abatement plans, including but not limited to those listed in Annex 1 of Attachment B' (to DCCEEW's RFI letter dated 16 July 2025).
- 2. When a native species or ecological community is listed as threatened under the EPBC Act, a Conservation Advice document is developed to assist its recovery. A Conservation Advice document guides recovery planning and identifies actions required for conservation and recovery of the threatened species or ecological community, and informs the Australian Government on required investments and regulatory decision-making.
- 3. Currently (August 2025) there are seven approved Conservation Advice documents that are directly relevant to the following seven key marine species in the CG area (web links):
 - Conservation Advice for <u>Orcaella heinsohni</u> (Australian Snubfin Dolphin), March 2025.
 - Conservation Advice for <u>Sousa sahulensis</u> (Australian Humpback Dolphin), March 2025.
 - Conservation Advice for <u>Glyphis glyphis</u> (Speartooth Shark), April 2014.
 - Conservation Advice for <u>Glyphis garricki</u> (Northern River Shark). April 2014.
 - Conservation Advice for <u>Pristis pristis</u> (Largetooth Sawfish), April 2014.
 - Conservation Advice for <u>Pristis zijsron</u> (Green Sawfish), undated.
 - Conservation Advice for <u>Pristis clavata</u> (Dwarf Sawfish), October 2009.
- 4. Apart from the two dolphin species, for which the Conservation Advice was developed in 2025, the advice for the two river shark species and three sawfish species date back over ten years to 2014 and 2009, and in some cases are based on even older data dating back to 2001, with very limited survey effort across northern Australia. More recent data based on more comprehensive, systematic surveys may well expand and refine the known geographical range and increase the population estimates for some species. An example is Kyne (2020), who recommends a down-listing of the Northern River Shark from 'endangered' to 'vulnerable', based on surveys that show a greater geographical range and larger population numbers than previously assessed.
- 5. Tables 2 to 8 present key elements of each of the Conservation Advice documents for the seven species listed above, and how these elements have been addressed by BKA for the CG marine sand proposal.

TABLE 2: Conservation Advice for Orcaella heinsohni (Australian Snubfin Dolphin)

In effect under EPBC Act from 5 March 2025.

Key element from the Conservation Advice



Image credit: I Beasley

Adult size: Up to 2.7 m.

1. Conservation status:

Australian Snubfin Dolphins are listed as <u>Vulnerable</u> under the EPBC Act and are also protected as both a Migratory species and as a cetacean (whales & dolphins), making the species a Matter of National Environmental Significance (MNES) under the EPBC Act.

How addressed by BKA

As an MNES species, BKA has given very high priority to assessing potential impacts of the proposal and developing relevant impact avoidance, mitigation and monitoring measures for this species, as presented in the following referral documents:

- EPBC Referral Report No. 2 - Setting & Existing Environment.

- Section 9 Marine Fauna describes marine fauna in the area based on all available information and the site surveys commissioned by BKA.
 - Section 9.4.1 Australian Snubfin Dolphin provides a specific description of this species in the area, based on all available information and the site surveys commissioned by BKA (see next item)

EPBC Referral Report No. 2 - Setting & Existing Environment - ANNEX 14 - Marine Fauna Surveys Report.

This presents the methods and results of marine fauna surveys commissioned by BKA, including for Snubfin Dolphins, including literature review, consultations with relevant experts and stakeholders, assessment of previous surveys conducted by others in the area (e.g Brown et al 2016 & 2017), and the dry- and wetseason surveys carried out in accordance with the National Guidelines for the Survey of Cetaceans, Marine Turtles and Dugong (DCCEEW 2024).

EPBC Referral Report No. 4 - Impact Assessments.

- Section 10 Impact Assessment Marine Fauna, assesses potential impacts of the proposal on marine fauna in accordance with WA EPA guidelines and impact mitigation hierarchy.
- Section 10.3.1 Snubfin & Humpback Dolphins applies the impact assessment to these species and finds no significant or residual impacts in accordance with WA EPA guidelines and impact mitigation hierarchy.

EPBC Referral Report No. 7 - Commonwealth Matters.

Section 10.3 - Specific Assessment for Snubfin Dolphins assesses
potential impacts of the proposal on this species in accordance with
the EPBC Act significant impact criteria and impact mitigation
hierarchy, and finds no significant or residual impacts in accordance
with these criteria.

- EPBC Supplementary Report No. 2 - Noise Assessment.

 This includes detailed modelling of predicated noise emissions from the SPV and assessment of potential auditory injury and behavioural impacts on Snubfin Dolphins, in accordance with the US NMFWS criteria (as required by WA EPA), using a risk assessment approach, and finds that potential impacts are negligible.

EPBC Supplementary Report No. 3 - Commonwealth Environmental Management Plan (C-EMP).

 This includes best practice impact avoidance, mitigation, monitoring and response actions for Snubfin Dolphins in accordance with the impact mitigation hierarchy, as summarized against Element 6 below.

2. Distribution & populations:

This species:

- is shy, cryptic and elusive, and will tend to move away from vessels and other human activity (unlike some other dolphin species that can be attracted to vessels, including to ride bow-waves),
- inhabits shallow, turbid, coastal, waters along coastline in the sub-tropical and tropical zones of Australia from Exmouth Gulf in the west to Brisbane in the east, and also coastal waters of

As reported in <u>EPBC Referral Report No. 2 - Setting & Existing Environment - ANNEX 14 - Marine Fauna Surveys Report</u>, the marine fauna surveys commissioned by BKA and previous surveys in the area by Brown et al (2016 & 2107) indicate that numbers of Snubfin Dolphins that utilize waters within CG itself are unlikely to be more than a few individuals (<10) to a few tens of individuals at most. These appear to be part of a larger population that also ranges outside of CG throughout the inner coastal waters of Joseph Bonaparte Gulf and along the coast both to the west and east of CG, where larger numbers were sighted by previous surveys.

During earlier surveys by Brown et al (2016, 2017) there was a total of 34 sightings over a nine-day survey period, noting that their survey area extended outside of CG into JBG and ~65 kms westwards along the coast to and up the Berkeley River, with many of their sightings being in these areas

Key element from the Conservation Advice How addressed by BKA southern Indonesia and Papua New Guinea outside of CG. They made no sightings in the Proposed Operational Area (see Figure 2), (POA) During BKA's dry-season survey in July 2023 there was a total of 11 has an estimated total population of <10.000 sightings, including two sightings in the POA, over an eight-day survey period mature individuals across its range in Northern covering 823 km of transects. Australia During BKA's wet-season survey in February 2024 there was a total of four subpopulations studied to date generally do not sightings, including two in the POA, over a nine-day survey period covering contain more than 150 mature individuals; and 850 km of transects. typically displays strong site fidelity to coastal It should be noted that separate sightings could be of the same individual(s), areas but has also shown evidence of so the actual number of individuals may be less than the number of sightings. connectivity and movement (observed high Positive photographic ID was only obtained for two separate individuals rates of temporary emigration) between local during the wet-season (Feb 2024) survey only, while Brown et al (2016, 2017) identified six distinct individuals (noting that their survey area included a much larger area outside of CG). The closest areas to CG with important populations as identified in the Conservation Advice are (Figure 2): Roebuck Bay at Broome (800 km west of CG), estimated population >130, Cygnet Bay in the West Kimberley region (600 km west of CG), estimated population of ~50; and Darwin Harbour and surrounding waters in the NT (400 km east of CG), estimated population up to 70. 3. Foraging behaviour, diet & critical habitat: This is consistent with the findings of EPBC Referral Report No. 2 - Setting & Existing Environment - ANNEX 14 - Marine Fauna Surveys Report, where most of the (very few sightings) within CG tended to be near and around The species is a generalist-opportunistic predator feeding on a wide variety of fish associated with Adolphus Island at the southern end of the main body of CG, and close to the shallow coastal and estuarine environments. shoreline where they feed. The Conservation Advice states that most of the The POA is located in deeper, open waters in the central part of CG, away prey identified within the stomachs of this species from coastal foraging areas. The sandy seabed within the POA, which is have been associated with shallow coastalhighly dynamic with constantly mobile sand-waves driven by extremely strong estuarine environments, suggesting feeding occurs tidal currents, does not provide suitable foraging habitat. near the coast and in river mouths. For the few sightings in the POA as listed against Element 2 above, the Habitat critical to the survival of the species is dolphins were swimming purposefully and directionally, indicating they were defined as shallow inshore coastal waters and transiting the open, deeper water of the POA, likely enroute between their estuarine habitats up to 10 km from a coastline preferred near-coast foraging areas. and/or 20 km from a freshwater outflow. Within this range, sites with a high density of teleost fish and cephalopods, such as mangroves and seagrass meadows, are considered important foraging habitat. No Critical Habitat as defined under section 207A of the EPBC Act has been identified or included in the Register of Critical Habitat. 4. Significance to First Nations people: Please refer: EPBC Referral Report No. 3 - Traditional Owner Matters. The Conservation Advice states that: EPBC Referral Report No. 6 - Consultation. cetaceans generally can be significant to the culture of coastal First Nations people; and BKA has undertaken and continues to undertake comprehensive in the CG and adjacent areas dolphins are consultations and engagement with the two First Nations groups in the CG known as yinga to the Balanggarra people and area, Balanggarra and Miriuwang-Gajerrong. Neither group has expressed are recognised as important marine species. concerns about Snubfin Dolphins and both groups have issued letters of support for the proposal, as presented in Annexes to EPBC Referral Report No. 3 - Traditional Owner Matters. 5. Main threats & potential impacts of the proposed The proposed operation will not cause loss and degradation of habitat from climate change, marine pollution or coastal development, bycatch and operation: entanglement in fishing gear or disease (pls refer section 3.2 above regarding The Conservation Advice lists the main threats to disease). Snubfin Dolphins as: The main potential impacts of the proposed operation on Snubfin Dolphins are potential vessel strike and potential impacts of underwater noise from the Sand Production Vessel (SPV)

Key element from the Conservation Advice How addressed by BKA loss and degradation of habitat from climate Vessel Strikes: change, marine pollution and coastal development. Potential vessel strikes are assessed in Section 10.3.1 - Snubfin & Humpback bycatch and entanglement in active fishing Dolphins of EPBC Referral Report No. 4 - Impact Assessments in accordance gear. with WA EPA guidelines and impact mitigation hierarchy, and in Section 10.3 disease. - Specific Assessment for Snubfin Dolphins of EPBC Referral Report No. 7 -- vessel interactions (vessel strikes); and Commonwealth Matters in accordance with the EPBC Act significant impact - anthropogenic underwater noise. criteria and impact mitigation hierarchy. The assessments find no significant or residual impacts in accordance with respective guidelines, criteria and the impact mitigation hierarchy, including through the application of impact prevention and mitigation measures. Potential vessel strikes will be prevented, mitigated and monitored as described in CEO 7 - Vessel Strikes in EPBC Supplementary Report No. 3 -Commonwealth Environmental Management Plan (C-EMP) as follows: CEO 7: Vessel Strikes: No significant negative impacts are caused to populations of surface-dwelling marine fauna in CG from vessel strikes by the SPV. Impact prevention: Very low likelihood of encounters due to: Very low occurrence of these species in the POA (as indicated by dedicated site surveys). Very low presence of the SPV in CG (1-2 days every 2 weeks with zero presence in CG for 86% of the time during the project Naturally elusive species: The 2 dolphin species found in CG (Snubfins & Humpbacks) are naturally shy and elusive, which unlike other dolphin species, avoid vessels. SPV Marine Fauna Observation & Avoidance (MFOA) measures (with TOs): Very low vessel speed: The SPV will operate at very low speeds (<2 knots) when loading sand in CG, allowing fauna to move away; and improving the effectiveness of MFOA measures. Impact mitigation: SPV MFOA measures (with TOs) (this is both an impact prevention & mitigation measure). Very low vessel speed: The SPV will operate at very low speeds (<2 knots) when loading sand in CG, allowing fauna to move away; and improving the effectiveness of MFOA measures (this is both an impact prevention & mitigation measure). Trigger Criteria (TCs), Trigger Response Actions (TRAs), Threshold Criteria (THCs), Threshold Contingency Actions (TCAs) and monitoring and reporting measures for CEO7 are specified in the C-EMP, in accordance with WA EPA criteria, which DCCEEW advised is the accepted template for this proposal. **Underwater Noise:** The design and construction of the SPV will include relevant best-practice noise reduction measures in accordance with the IMO Underwater Radiated Noise Guidelines (IMO 2023), which constitute the international standard. Potential impacts of underwater noise are assessed in **EPBC Supplementary** Report No. 2 - Noise Assessment. This includes detailed modelling of predicated noise emissions from the SPV and assessment of potential auditory injury and behavioural impacts on Snubfin Dolphins, in accordance with the US NMFWS criteria (as required by WA EPA), using a risk assessment approach, and finds that potential impacts are negligible. Despite the assessment that potential impacts are negligible, as a precautionary measure BKA proposes to undertake monitoring of underwater noise in CG during commencement of operations to assess compliance with the findings of the Noise Assessment, as described in EPBC Supplementary Report No. 3 - Commonwealth Environmental Management Plan (C-EMP).

Key element from the Conservation Advice How addressed by BKA 6. Survey & monitoring priorities: BKA commissioned appropriate baseline surveys in accordance with the National Guidelines for the Survey of Cetaceans, Marine Turtles and Dugong, as reported in <u>EPBC Referral Report No. 2 - Setting & Existing Environment - ANNEX 14 - Marine Fauna Surveys Report</u>. This included: The Conservation Advice lists a number of survey and monitoring priorities, which are strategic in nature for adoption by relevant parties involved in literature search and review, the conservation of the species at the national consultations with relevant experts and stakeholders, level, and are not targeted at specific development assessment of previous surveys conducted by others in the area (e.g proposals. The following are of direct relevance to Brown et al 2016 & 2017), the proposal: an eight-day dry-season survey carried out in July 2023, covering 823 km of transects, undertake appropriate baseline surveys in a nine-day wet-season survey carried out in February 2024, covering 850 accordance with the National Guidelines for km of transects: and the Survey of Cetaceans, Marine Turtles and 49 days of incidental marine fauna observations during other Dugong (DCCEEW 2024), environmental survey work in CG, in both the dry- and wet-seasons. undertake continued monitoring; and standardised monitoring in collaboration with BKA proposes to undertake continued monitoring of Snubfin Dolphins during First Nations Sea Ranger groups. the project period, through the Marine Fauna Observation and Avoidance (MFOA) program described in EPBC Supplementary Report No. 3 -Commonwealth Environmental Management Plan (C-EMP). BKA proposes to contract, fund, train and equip the local First Nations rangers to undertake the proposed MFOA program, and discussions on this have been included in BKA's consultations with the First Nations groups to 7. Information & research priorities: Underwater noise: The design and construction of the SPV will include relevant best-practice The Conservation Advice lists a number of noise reduction measures in accordance with the IMO Underwater Radiated information and research priorities, which are Noise Guidelines (IMO 2023), which constitute the international standard. strategic in nature for adoption by relevant parties involved in the conservation of the species at the national level, and are not targeted at specific Potential impacts of underwater noise are assessed in EPBC Supplementary development proposals. The following are of direct Report No. 2 - Noise Assessment. This includes detailed modelling of relevance to the proposal: predicated noise emissions from the SPV and assessment of potential auditory injury and behavioural impacts on Snubfin Dolphins, in accordance with the US NMFWS criteria (as required by WA EPA), using a risk Underwater anthropogenic noise: Further assessment approach, and finds that potential impacts are negligible. understand the impact of noise pollution from increasing vessel traffic on dolphin Despite the assessment that potential impacts are negligible, as a distribution and behaviour. precautionary measure BKA proposes to undertake monitoring of underwater noise in CG during commencement of operations to assess compliance with Vessel interactions: the findings of the Noise Assessment, as described in EPBC Supplementary ensure the risk of vessel strike is Report No. 3 - Commonwealth Environmental Management Plan (C-EMP). considered when assessing actions that increase vessel traffic in areas where Vessel interactions: the species occurs, and, if required, implement appropriate mitigation As outlined in the responses to Elements 5 and 6 above, the issue of vessel measures; and interactions is comprehensively addressed, including a MFOA program and ensure all vessel strike incidents are reporting via the National Ship Strike Database. reported in the National Ship Strike Database 8. Recovery Plan: The measures to be implemented by BKA as summarized in this table, including the data that will be generated by the proposed monitoring program, will contribute to improved understanding of the species and to their The Conservation Advice states that the Threatened Species Scientific Committee conservation and recovery. recommended that a Recovery Plan is not required for this species. The Conservation Advice itself provides sufficient guidance for implementing priority conservation actions, mitigating key threats and supporting recovery.




FIGURE 2: Distribution of Australian Snubfin Dolphin in Australian waters (also found in southern Indonesia & PNG) (source: DCCEEW)

TABLE 3: Conservation Advice for Sousa sahulensis (Australian Humpback Dolphin)

In effect under EPBC Act from 5 March 2025.

Key element from the Conservation Advice

Image credit: A Brown

Adult size: Up to 2.7 m.

1. Conservation status:

Australian Humpback Dolphins are listed as <u>Vulnerable</u> under the EPBC Act and are also protected as both a Migratory species and as a cetacean (whales & dolphins), making the species a Matter of National Environmental Significance (MNES) under the EPBC Act.

How addressed by BKA

As an MNES species, BKA has given very high priority to assessing potential impacts of the proposal and developing relevant impact avoidance, mitigation and monitoring measures for this species, as presented in the following referral documents:

- EPBC Referral Report No. 2 - Setting & Existing Environment.

- Section 9 Marine Fauna describes marine fauna in the area based on all available information and the site surveys commissioned by BKA.
- Section 9.4.2 Australian Humpback Dolphin provides a specific description of this species in the area, based on all available information and the site surveys commissioned by BKA (see next item)

- <u>EPBC Referral Report No. 2 - Setting & Existing Environment - ANNEX 14 - Marine Fauna Surveys Report.</u>

This presents the methods and results of marine fauna surveys commissioned by BKA, including for Humpback Dolphins, including literature review, consultations with relevant experts and stakeholders, assessment of previous surveys conducted by others in the area (e.g Brown et al 2016 & 2017), and the dry- and wetseason surveys carried out in accordance with the National Guidelines for the Survey of Cetaceans, Marine Turtles and Dugong (DCCEEW 2024).

EPBC Referral Report No. 4 - Impact Assessments.

- Section 10 Impact Assessment Marine Fauna, assesses potential impacts of the proposal on marine fauna in accordance with WA EPA guidelines and impact mitigation hierarchy.
- Section 10.3.1 Snubfin & Humpback Dolphins applies the impact assessment to these species and finds no significant or residual impacts in accordance with WA EPA guidelines and impact mitigation hierarchy.

EPBC Referral Report No. 7 - Commonwealth Matters.

Section 10.4 - Specific Assessment for Snubfin Dolphins assesses
potential impacts of the proposal on this species in accordance with
the EPBC Act significant impact criteria and impact mitigation
hierarchy, finds no significant or residual impacts in accordance with
these criteria, and applies equally to Humpback Dolphins.

- EPBC Supplementary Report No. 2 - Noise Assessment.

 This includes detailed modelling of predicated noise emissions from the SPV and assessment of potential auditory injury and behavioural impacts on Humpback Dolphins, in accordance with the US NMFWS criteria (as required by WA EPA), using a risk assessment approach, and finds that potential impacts are negligible.

EPBC Supplementary Report No. 3 - Commonwealth Environmental Management Plan (C-EMP).

 This includes best practice impact avoidance, mitigation, monitoring and response actions for Humpback Dolphins in accordance with the impact mitigation hierarchy, as summarized against Element 6 below.

2. Distribution & populations:

This species:

- is shy, cryptic and elusive, and will tend to move away from vessels and other human activity (unlike some other dolphin species that can be attracted to vessels, including to ride bowwaves).
- inhabits shallow, turbid, coastal waters along coastline in the sub-tropical and tropical zones of Australia from Shark Bay in the west to the Queensland / NSW border area in the east, and

As reported in EPBC Referral Report No. 2 - Setting & Existing Environment - ANNEX 14 - Marine Fauna Surveys Report, the marine fauna surveys commissioned by BKA and previous surveys in the area by Brown et al (2016 & 2107) indicate that numbers of Humpback Dolphins that utilize waters within CG itself are unlikely to be more than a few individuals (<10) to a few tens of individuals at most. These appear to be part of a larger population that also ranges outside of CG throughout the inner coastal waters of Joseph Bonaparte Gulf and along the coast both to the west and east of CG, where larger numbers were sighted by previous surveys.

During earlier surveys by Brown et al (2016, 2017) there was a total of 42 sightings over a nine-day survey period, noting that their survey area extended outside of CG into JBG and ~65 kms westwards along the coast to and up the Berkeley River. Most Humpback Dolphin sightings where near

Key element from the Conservation Advice also coastal waters of southern Indonesia and Papua New Guinea (see Figure 3), has an estimated total population of <10.000

- mature individuals across its range in Northern Australia
- subpopulations studied to date generally do not contain more than 150 mature individuals; and
- typically displays strong site fidelity to coastal areas but has also shown evidence of connectivity and movement (observed high rates of temporary emigration) between local sites.

Surveys have highlighted the importance of riverineestuarine systems to the species with a number of populations inhabiting river mouths, tidal rivers and estuaries across Northern Australia.

The closest areas to CG with important populations as identified in the Conservation Advice are (Figure 3):

- North West Cape to Pilbara region (1,500 km west of CG), estimated population up to 2,910,
- Cygnet Bay in the West Kimberley region (600 km west of CG), estimated population up to 20;
- Darwin Harbour and surrounding waters in the NT (400 km east of CG), estimated population up to 99.

How addressed by BKA

Cape Dussejour to the west of CG and outside CG and along the coast to the west. They made no sightings in the POA.

During BKA's dry-season survey in July 2023 there were no sightings of Humpback Dolphins, over an eight-day survey period covering 823 km of transects

During BKA's wet-season survey in February 2024 there was a single, unconfirmed sighting of a possible Humpback Dolphin just to the north of the POA, towards Cape Dussejour, over a nine-day survey period covering 850 km of transects

It should be noted that separate sightings could be of the same individual(s), so the actual number of individuals may be less than the number of sightings. Brown et al (2016, 2017) identified 12 distinct individuals (noting that their survey area included a much a larger area outside of CG).

3. Foraging behaviour, diet & critical habitat:

The species is a generalist-opportunistic predator feeding on a wide variety of fish associated with shallow coastal and estuarine environments.

The species has been observed feeding in inshore coastal and estuarine habitats such as rivers and creeks, on exposed inter-tidal banks and flats and over seagrass meadows, rocks and reef.

Habitat critical to the survival of the species is defined as shallow inshore coastal waters and estuarine habitats up to 20 km from a coastline or land body, such as an island group, with sand banks, mud flats, seagrass, rock and/or reef substrate. Within this range, sites with a high density of teleost fish, cephalopods and bivalves are important foraging habitat.

No Critical Habitat as defined under section 207A of the EPBC Act has been identified or included in the Register of Critical Habitat.

This is consistent with the findings of EPBC Referral Report No. 2 - Setting & Existing Environment - ANNEX 14 - Marine Fauna Surveys Report, where most of the (very few) sightings were near near Cape Dussejour to the west to CG and outside and along the coast to the west of CG, and none within

There is an area of expansive inter-tidal banks along the coast just south of Cape Dussejour, and Humpback Dolphins are known to target such areas for feeding

The POA is located in deeper, open waters in the central part of CG, away from coastal foraging areas. The sandy seabed within the POA, which is highly dynamic with constantly mobile sand-waves driven by strong tidal currents, does not provide suitable foraging habitat.

4. Significance to First Nations people:

The Conservation Advice states that:

- cetaceans generally can be significant to the culture of coastal First Nations people; and
- in the CG and adjacent areas dolphins are known as yinga to the Balanggarra people and are recognised as important marine species.

- EPBC Referral Report No. 3 Traditional Owner Matters.
- EPBC Referral Report No. 6 Consultation.

BKA has undertaken and continues to undertake comprehensive consultations and engagement with the two First Nations groups in the CG area, Balanggarra and Miriuwang-Gajerrong. Neither group has expressed concerns about Humpback Dolphins and both groups have issued letters of support for the proposal, as presented in Annexes to EPBC Referral Report No. 3 - Traditional Owner Matters.

5. Main threats & potential impacts of the proposed operation:

The Conservation Advice lists the main threats to Snubfin Dolphins as

The proposed operation will not cause loss and degradation of habitat from climate change, marine pollution or coastal development, bycatch and entanglement in fishing gear or disease (pls refer section 3.2 above regarding disease).

Key element from the Conservation Advice

- loss and degradation of habitat from climate change, marine pollution and coastal development,
- bycatch and entanglement in active fishing gear,
- disease,
- vessel interactions (vessel strikes); and
- anthropogenic underwater noise.

How addressed by BKA

The main potential impacts of the proposed operation on Humpback Dolphins are potential vessel strike and potential impacts of underwater noise from the Sand Production Vessel (SPV).

Potential vessel strikes are assessed in Section 10.3.1 - Snubfin & Humpback Dolphins of EPBC Referral Report No. 4 - Impact Assessments in accordance with WA EPA guidelines and impact mitigation hierarchy, and in Section 10.3 - Specific Assessment for Snubfin Dolphins of EPBC Referral Report No. 7 - Commonwealth Matters in accordance with the EPBC Act significant impact criteria and impact mitigation hierarchy (which also applies to Humpback Dolphins). The assessments find no significant or residual impacts in accordance with respective guidelines, criteria and the impact mitigation hierarchy, including through the application of impact prevention and mitigation measures.

Potential vessel strikes will be prevented, mitigated and monitored as described in CEO 7 - Vessel Strikes in <u>EPBC Supplementary Report No. 3 - Commonwealth Environmental Management Plan (C-EMP)</u>, as outlined for Snubfin Dolphins in Table 2 above, which is not repeated here for reasons of economy.

Potential impacts of underwater noise are addressed as outlined for Snubfin Dolphins in Table 2 above, which is not repeated here for reasons of economy.

6. Survey & monitoring priorities:

The Conservation Advice lists a number of survey and monitoring priorities, which are strategic in nature for adoption by relevant parties involved in the conservation of the species at the national level, and are not targeted at specific development proposals. The following are of direct relevance to the proposal:

- undertake appropriate baseline surveys in accordance with the National Guidelines for the Survey of Cetaceans, Marine Turtles and Dugong (DCCEEW 2024),
- undertake continued monitoring; and
- standardised monitoring in collaboration with First Nations Sea Ranger groups.

BKA commissioned appropriate baseline surveys in accordance with the National Guidelines for the Survey of Cetaceans, Marine Turtles and Dugong, as reported in Environment - ANNEX 14 - Marine Fauna Surveys Report. This included:

- literature search and review,
 consultations with relevant experts and stakeholders,
- assessment of previous surveys conducted by others in the area (e.g Brown et al 2016 & 2017).
- an eight-day dry-season survey carried out in July 2023, covering over 820 km of transects,
- a nine-day wet-season survey carried out in February 2024, covering over 850 km of transects; and
- 49 days of incidental marine fauna observations during other environmental survey work in CG, in both the dry- and wet-seasons.

BKA proposes to undertake continued monitoring of Humpback Dolphins during the project period, through the Marine Fauna Observation and Avoidance (MFOA) program described in EPBC Supplementary Report No.3 - Commonwealth Environmental Management Plan (C-EMP).

BKA proposes to contract, fund, train and equip the local First Nations rangers to undertake the proposed MFOA program, and discussions on this have been included in BKA's consultations with the First Nations groups to date.

7. Information & research priorities:

The Conservation Advice lists a number of information and research priorities, which are strategic in nature for adoption by relevant parties involved in the conservation of the species at the national level, and are not targeted at specific development proposals. The following are of direct relevance to the proposal:

- Underwater anthropogenic noise: Further understand the impact of noise pollution from increasing vessel traffic on dolphin distribution and behaviour.
- Vessel interactions:
 - ensure the risk of vessel strike is considered when assessing actions that increase vessel traffic in areas where the species occurs, and, if required, implement appropriate mitigation measures; and

Underwater Noise:

The design and construction of the SPV will include relevant best-practice noise reduction measures in accordance with the *IMO Underwater Radiated Noise Guidelines* (IMO 2023), which constitute the international standard.

Potential impacts of underwater noise are assessed in EPBC Supplementary Report No. 2 - Noise Assessment. This includes detailed modelling of predicated noise emissions from the SPV and assessment of potential auditory injury and behavioural impacts on Humpback Dolphins, in accordance with the US NMFWS criteria (as required by WA EPA), using a risk assessment approach, and finds that potential impacts are negligible.

Despite the assessment that potential impacts are negligible, BKA proposes to undertake monitoring of underwater noise in CG during commencement of operations to assess compliance with the findings of the Noise Assessment, as described in Environmental Management Plan (C-EMP).

Vessel interactions:

As outlined in the responses to Elements 5 and 6 above, the issue of vessel interactions is comprehensively addressed, including a MFOA program and reporting via the National Ship Strike Database.

Key element from the Conservation Advice	How addressed by BKA
 ensure all vessel strike incidents are reported in the National Ship Strike Database. 	
8. Recovery Plan: The Conservation Advice states that the Threatened Species Scientific Committee recommended that a Recovery Plan is not required for this species. The Conservation Advice itself provides sufficient guidance for implementing priority conservation actions, mitigating key threats and supporting	The measures to be implemented by BKA as summarized in this table, including the data that will be generated by the proposed monitoring program, will contribute to improved understanding of the species and to their conservation and recovery.




FIGURE 3: Distribution of Australian Humpback Dolphin in Australian waters (also found in southern Indonesia & PNG) (source: DCCEEW)

TABLE 4: Conservation Advice for Glyphis glyphis (Speartooth Shark)

In effect under EPBC Act from 11 April 2014.

Key element from the Conservation Advice

Image credit: Marinewise

Adult size: Up to 2 m.

1. Conservation Status:

Speartooth Sharks are listed as <u>Critically Endangered</u> under the EPBC Act, making the species a Matter of National Environmental Significance (MNES) under that Act.

NOTE:

- The Conservation Advice is dated 11 April 2014 and states that the Conservation Status is based on limited data from nearly 25 years ago in 2001, which indicated a limited geographical distribution and low population numbers, but with significant gaps in survey effort across Northern Australia.
- Re-assessment using more recent data may likely suggest a down-listing of this species, as recommended by Kyne (2020) for the closely related Northern River Shark (*Glyphis garricki*) (see Table 5 below).

How addressed by BKA

As an MNES species, BKA has given very high priority to assessing potential impacts of the proposal and developing relevant impact avoidance, mitigation and monitoring measures for this species, as presented in the following referral documents:

- EPBC Referral Report No. 2 - Setting & Existing Environment.

- Section 9 Marine Fauna describes marine fauna in the area based on all available information and the site surveys commissioned by BKA
- Section 9.4.6 River Sharks provides a specific description of this species in the area, based on all available information and the eDNA site surveys commissioned by BKA (see below).

EPBC Referral Report No. 2 - Setting & Existing Environment - ANNEX 13 - Marine eDNA Report.

- This presents the methods and results of marine eDNA surveys commissioned by BKA and undertaken by the University of Canberra National eDNA Reference Centre.
- The surveys collected and analysed 60 seabed sediment samples and 26 water samples from across 20 separate sites within the POA, in other open-water parts of CG and up the inlets, creeks and rivers on both the eastern and western sides of CG (but not as far upstream as the Lower Ord River due to the long distance from the POA - > 35 km).
- No eDNA evidence of Speartooth Sharks was identified by this this survey.

EPBC Referral Report No. 2 - Setting & Existing Environment - ANNEX 14 -Marine Fauna Surveys Report.

This presents the methods and results of marine fauna surveys commissioned by BKA, including for any shark species, including literature review, consultations with relevant experts and stakeholders, assessment of previous surveys conducted by others in the area (e.g Kyne et al 2020), and the dry- and wet-season surveys carried out in accordance with the National Guidelines for the Survey of Cetaceans, Marine Turtles and Dugong (DCCEEW 2024). No Speartooth Sharks were observed during these surveys.

EPBC Referral Report No. 4 - Impact Assessments.

- Section 10 Impact Assessment Marine Fauna, assesses potential impacts of the proposal on marine fauna in accordance with WA EPA guidelines and impact mitigation hierarchy.
- Section 10.3.6 River Sharks applies the impact assessment to these species and finds no significant or residual impacts in accordance with WA EPA guidelines and impact mitigation hierarchy.

EPBC Referral Report No. 7 - Commonwealth Matters.

 Section 10.4 assesses potential impacts of the proposal on listed species including river sharks in accordance with the EPBC Act significant impact criteria and impact mitigation hierarchy, and finds no significant or residual impacts in accordance with these criteria.

EPBC Supplementary Report No. 3 - Commonwealth Environmental Management Plan (C-EMP).

 This includes best practice impact avoidance, mitigation, monitoring and response actions for river sharks in accordance with the impact mitigation hierarchy, as summarized against Element 6 below.

2. Distribution & populations:

The Conservation Advice states that:

 Speartooth sharks are capable of living in and moving between freshwater and seawater. Juveniles and sub-adults utilise large tropical mangrove-lined river systems with lower salinities as their primary habitat, often being found well upstream, including in near-fresh waters. As reported in Section 9.4.6 of <u>EPBC Referral Report No. 2 - Setting & Existing Environment</u>, Kyne et al (2021) reported sampling juvenile Speartooth Sharks in the Lower Ord River ~35 km upstream from the main body of CG, in 2015 and 2019, consistent with their preference for less saline, upstream waters of rivers and estuaries.

There are no records of this species in the more saline, deeper marine waters of the main body of CG where the POA is located. As outlined above, the eDNA sampling commissioned by BKA did not identify evidence of Speartooth Sharks. However, the occasional adult could potentially pass through that area during inshore/offshore movements.

Key element from the Conservation Advice	How addressed by BKA
Individuals have a tidally influenced movement pattern, moving up and downstream with the flood and ebb tides, and primarily swim well above the seabed. Surveys show that individuals repeatedly utilise small sections of the available habitat.	
Based on physiological and life history similarities with Bull Sharks (<i>Carcharhinus leucas</i>), it is assumed that adult Speartooth Sharks may live outside of rivers in the coastal marine environment.	
Their currently known geographical range covers the tropical river systems and coastal waters from the Kimberley region of WA to the east coast of tropical Queensland and rivers along the southern coast of PNG. The Conservation Advice (based on old 2001 data) states that there have been no confirmed records from Queensland's east coast since 1983, indicating that they may have become locally extinct in that area (possibly due to historically extensive commercial gill-netting for Barramundi and other finfish species).	
Knowledge of their overall distribution is constrained by a lack of surveys across Northern Australia. Increased survey effort may likely expand their known geographical range, as reported by Kyne (2020) for the closely related Northern River Shark (Glyphis garricki) (see Table 5 below).	
 Knowledge of their population structure and numbers is also constrained by a lack of surveys across Northern Australia, and there is currently no overall population estimate available. 	
3. Foraging behaviour, diet & critical habitat: The Conservation Advice does not contain information on the foraging behaviour and diet of this species. As a Carcharhinid shark it would actively hunt and opportunistically prey upon a wide variety of smaller marine species throughout the water column. As outlined above the habitat for this species is tropical mangrove-lined river systems and estuaries for juveniles and sub-adults, and is likely be the coastal waters offshore from these river and estuarine areas for adults, although very little data is available for the latter. The Conservation Advice states that the distribution of this species is not known to overlap with any EPBC Act-listed threatened ecological communities.	The main habitat for juvenile and sub-adult Speartooth Sharks in CG is located in the Lower Ord River ~35 km upstream from the POA. There is therefore no overlap of the proposed operation with this habitat or potential for direct impacts on juveniles and sub-adults. As outlined above there are no records of this species in the more saline, deeper marine waters of the main body of CG where the POA is located, although the occasional adult could potentially pass through that area during inshore/offshore movements. Potential impacts on sharks moving through the POA, and proposed impact prevention, mitigation and monitoring measures are summarised against Element 5 below.
4. Significance to First Nations people: The Conservation Advice states that: Fishing of sharks and rays is a part of traditional fishing practices and historically makes up an important part of the diet of coastal indigenous communities. Indigenous Australians are allowed to take and eat Speartooth Sharks for personal, domestic or non-commercial communal needs.	Please refer: - EPBC Referral Report No. 3 - Traditional Owner Matters. - EPBC Referral Report No. 6 - Consultation. BKA has undertaken and continues to undertake comprehensive consultations and engagement with the two First Nations groups in the CG area, Balanggarra and Miriuwang-Gajerrong. Neither group has expressed concerns about shark species and both groups have issued letters of support for the proposal, as presented in Annexes to EPBC Referral Report No. 3 - Traditional Owner Matters.

Key element from the Conservation Advice	How addressed by BKA
S. Main threats & potential impacts of the proposed operation: The Conservation Advice lists the main threats to Speartooth Sharks as: commercial fishing (especially gill netting and prawn trawling), recreational fishing, indigenous fishing, illegal, unreported and unregulated (IUU) fishing, entanglement in marine debris; and habitat degradation and modification.	The proposed operation will not contribute to any of the threats listed in the Conservation Advice. It does not include any form of fishing, it will not discharge debris into the marine environment (see section 3.5 below) and it will not degrade or modify the species' habitat. As outlined above the occasional adult could potentially pass through the POA during inshore/offshore movements, which would present a very low potential for vessel strike by the SPV, or a negligible risk of being entrained in the SPV's drag-head when it is operating. Factors and measures to prevent, mitigate, monitor and respond to potential vessel strikes are the same as those described for both Snubfin and Humpback Dolphins against Element 5 in Tables 2 and 3 above, and are not repeated here for reasons of economy. The potential for this species to be entrained in the SPV's drag-head when it is operating is negligible for the following reasons: Apparent absence of this species in the POA (based on surveys and eDNA sampling). Very low presence of the SPV in CG (1-2 days every 2 weeks with zero presence in CG for 86% of the time during the project lifespan).
	 Very low vessel speed: The SPV will operate at very low speeds (<2 knots) when loading sand in CG, allowing fauna to move away; and improving the effectiveness of MFOA measures. Swimming behaviour: The fact that this species primarily swims well above the seabed (the drag-head operates on the seabed). Single drag-head: The SPV will only have a single suction-arm and draghead (standard TSHD's usually have a pair of suction-arms and dragheads). Marine fauna excluder ('turtle tickler chains'): Will be fitted in front of the drag-head. The potential entrainment of a river shark in the drag-head would not constitute significant impact on the species stock or population as defined by
6. Research Priorities & Priority Actions: The Conservation Advice lists a number of research priorities and priority actions, which are strategic in nature for adoption by relevant parties involved in the conservation of the species at the national level. The research priorities and priority actions are designed primarily to address the key threats outlined against Element 5 above.	The proposed operation will not contribute to any of the threats listed in the Conservation Advice which are addressed by the research priorities and priority actions.
7. Recovery Plan: The Conservation Advice references a Draft Sawfish and River Sharks Multispecies Recovery Plan, which was published as a final plan in 2015.	Please refer Table 9 in section 3.4 below which presents BKA's response to relevant elements of the Sawfish and River Sharks Multispecies Recovery Plan.

TABLE 5: Conservation Advice for Glyphis garricki (Northern River Shark)

In effect under EPBC Act from 11 April 2014.

Key element from the Conservation Advice

Image credit: Sam Lyne

Adult size: Up to 2.5 m.

1. Conservation Status:

Northern River Sharks are listed as Endangered under the EPBC Act, making the species a Matter of National Environmental Significance (MNES) under that Act

NOTE:

- The Conservation Advice is dated 11 April 2014 and states that the Conservation Status is based on limited data from 2001 which indicated a limited geographical distribution and low population numbers.
- Kyne (2020) reported sampling for Northern River Sharks in 11 rivers in the NT and WA, starting in 2013. The species was found in the Lower Ord, Durack and Pentecost Rivers upstream from CG. The report estimates the total Australian population size to be between 2,500 and 10,000 adults, and recommends a down-listing of this species from 'endangered' to 'vulnerable'.

How addressed by BKA

As an MNES species, BKA has given very high priority to assessing potential impacts of the proposal and developing relevant impact avoidance, mitigation and monitoring measures for this species, as presented for the Element in Table 4 for Speartooth Sharks above, which is not repeated here for reasons of economy.

2. Distribution & populations:

The Conservation Advice states that:

- Northern River Sharks have a similar distribution and populations to Speartooth Sharks as described in Table 4 above, including living in and moving between freshwater and seawater. As for Speartooth Sharks, juveniles and sub-adults utilise large tropical mangrove-lined river systems with lower salinities as their primary habitat, often being found well upstream, including in near-fresh waters, while adults may live outside of rivers in the coastal marine environment.
- Their currently known geographical range covers the tropical river systems and coastal waters from the Kimberley region of WA to the NT side of the Gulf of Carpentaria and rivers along the southern coast of PNG. The Conservation Advice does not mention Queensland waters, although this may simply reflect a lack of survey coverage.
- Increased survey effort may likely expand their known geographical range, as reported by Kyne (2020).
- Knowledge of their population structure and numbers is also constrained by a lack of surveys across Northern Australia. As outlined above, Kyne (2020) estimates the total Australian population size to be

As reported in Section 9.4.6 of <u>EPBC Referral Report No. 2 - Setting & Existing Environment</u>. Kyne et al (2021) reported sampling juvenile Northern River Sharks in the Lower Ord River ~35 km upstream from the main body of CG, and in the Durack and Pentecost Rivers >80 km upstream from CG, in 2015 and 2019, consistent with their preference for less saline, upstream waters of rivers and estuaries.

There are no records of this species in the more saline, deeper marine waters of the main body of CG where the POA is located. As outlined above, the eDNA sampling commissioned by BKA did not identify evidence of Speartooth Sharks. However, the occasional adult could potentially pass through that area during inshore/offshore movements.

Key element from the Conservation Advice	How addressed by BKA
between 2,500 and 10,000 adults, although he does not seem to have undertaken surveys in Queensland waters.	
3. Foraging behaviour, diet & critical habitat: The Conservation Advice does not contain information on the foraging behaviour and diet of this species. As a Carcharhinid shark it would actively hunt and opportunistically prey upon a wide variety of smaller marine species throughout the water column. As outlined above the habitat for this species is tropical mangrove-lined river systems and estuaries for juveniles and sub-adults, and is likely be the coastal waters offshore from these river and estuarine areas for adults, although very little data is available for the latter. The Conservation Advice states that the distribution of this species is not known to overlap with any EPBC Act-listed threatened ecological communities.	The main habitat for juvenile and sub-adult Northern River Sharks in CG is located in the Lower Ord River ~35 km upstream from the POA, and in the Durack and Pentecost Rivers >80 km upstream from CG. There is therefore no overlap of the proposed operation with this habitat or potential for direct impacts on juveniles and sub-adults. As outlined above there are no records of this species in the more saline, deeper marine waters of the main body of CG where the POA is located, although the occasional adult could potentially pass through that area during inshore/offshore movements. Potential impacts on sharks moving through the POA, and proposed impact prevention, mitigation and monitoring measures are summarised against Element 5 below.
4. Significance to First Nations people: The Conservation Advice states that: Fishing of sharks and rays is a part of traditional fishing practices and historically makes up an important part of the diet of coastal indigenous communities. Indigenous Australians are allowed to take and eat Speartooth Sharks for personal, domestic or non-commercial communal needs.	Please refer: - EPBC Referral Report No. 3 - Traditional Owner Matters. - EPBC Referral Report No. 6 - Consultation. BKA has undertaken and continues to undertake comprehensive consultations and engagement with the two First Nations groups in the CG area, Balanggarra and Miriuwang-Gajerrong. Neither group has expressed concerns about shark species and both groups have issued letters of support for the proposal, as presented in Annexes to EPBC Referral Report No. 3 - Traditional Owner Matters.
5. Main threats & potential impacts of the proposed operation: The Conservation Advice lists the main threats to Speartooth Sharks as: - commercial fishing (especially gill netting and prawn trawling), - recreational fishing, - indigenous fishing, - illegal, unreported and unregulated (IUU) fishing, - entanglement in marine debris; and - habitat degradation and modification.	The proposed operation will not contribute to any of the threats listed in the Conservation Advice. It does not include any form of fishing, it will not discharge debris into the marine environment (see section 3.5 below) and it will not degrade or modify the species' habitat. As outlined above the occasional adult could potentially pass through the POA during inshore/offshore movements, which would present a very low potential for vessel strike by the SPV, or a negligible risk of being entrained in the SPV's drag-head when it is operating. Factors and measures to prevent, mitigate, monitor and respond to potential vessel strikes are the same as those described for both Snubfin and Humpback Dolphins against Element 5 in Tables 2 and 3 above, and are not repeated here for reasons of economy. The potential for this species to be entrained in the SPV's drag-head when it is operating is negligible for the following reasons: Apparent absence of this species in the POA (based on surveys and eDNA sampling). Very low presence of the SPV in CG (1-2 days every 2 weeks with zero presence in CG for 86% of the time during the project lifespan). Very low vessel speed: The SPV will operate at very low speeds (<2 knots) when loading sand in CG, allowing fauna to move away; and improving the effectiveness of MFOA measures. Swimming behaviour: The fact that this species primarily swims well above the seabed (the drag-head operates on the seabed). Single drag-head: The SPV will only have a single suction-arm and drag-head (standard TSHD's usually have a pair of suction-arms and drag-heads). Marine fauna excluder ('turtle tickler chains'): Will be fitted in front of the drag-head. The potential entrainment of a river shark in the drag-head would not constitute significant impact on the species stock or population as defined by the EPBC Act significant impact on the species stock or population as defined by the EPBC Act significant impact criteria.

Key element from the Conservation Advice	How addressed by BKA
6. Research Priorities & Priority Actions: The Conservation Advice lists a number of research priorities and priority actions, which are strategic in nature for adoption by relevant parties involved in the conservation of the species at the national level. The research priorities and priority actions are designed primarily to address the key threats outlined against Element 5 above.	The proposed operation will not contribute to any of the threats listed in the Conservation Advice which are addressed by the research priorities and priority actions.
7. Recovery Plan: The Conservation Advice references a Draft Sawfish and River Sharks Multispecies Recovery Plan, which was published as a final plan in 2015.	Please refer Table 9 in section 3.4 below which presents BKA's response to relevant elements of the Sawfish and River Sharks Multispecies Recovery Plan.

TABLE 6: Conservation Advice for <u>Pristis pristis</u> (Largetooth Sawfish) (also known as Freshwater Sawfish) In effect under EPBC Act from 11 April 2014.

Key element from the Conservation Advice

Image credit: Fishes of Aus

Adult size: Up to 6.5 m.

1. Conservation Status:

Largetooth Sawfish are listed as <u>Vulnerable</u> under the EPBC Act, making the species a Matter of National Environmental Significance (MNES) under that Act.

How addressed by BKA

As an MNES species BKA has given very high priority to assessing potential impacts of the proposal and developing relevant impact avoidance, mitigation and monitoring measures for this species, as presented in the following referral documents:

- EPBC Referral Report No. 2 - Setting & Existing Environment.

- Section 9 Marine Fauna describes marine fauna in the area based on all available information and the site surveys commissioned by RKΔ
- Section 9.4.5 Sawfish provides a specific description of this species in the area, based on all available information and the eDNA site surveys commissioned by BKA (see below).

EPBC Referral Report No. 2 - Setting & Existing Environment - ANNEX 13 - Marine eDNA Report.

- This presents the methods and results of marine eDNA surveys commissioned by BKA and undertaken by the University of Canberra National eDNA Reference Centre.
- The surveys collected and analysed 60 seabed sediment samples and 26 water samples from across 20 separate sites within the POA, in other open-water parts of CG and up the inlets, creeks and rivers on both the eastern and western sides of CG (but not as far upstream as the Lower Ord River due to the long distance from the POA - > 35 km).
- No eDNA evidence of Largetooth Sawfish was identified by this this survey.

EPBC Referral Report No. 2 - Setting & Existing Environment - ANNEX 14 -Marine Fauna Surveys Report.

This presents the methods and results of marine fauna surveys commissioned by BKA, including for any shark species, including literature review, consultations with relevant experts and stakeholders, assessment of previous surveys conducted by others in the area (e.g Kyne et al 2020), and the dry- and wet-season surveys carried out in accordance with the National Guidelines for the Survey of Cetaceans, Marine Turtles and Dugong (DCCEEW 2024). No Largetooth Sawfish were observed during these surveys.

EPBC Referral Report No. 4 - Impact Assessments.

- Section 10 Impact Assessment Marine Fauna, assesses potential impacts of the proposal on marine fauna in accordance with WA EPA guidelines and impact mitigation hierarchy.
- Section 10.3.5 Sawfish applies the impact assessment to these species and finds no significant or residual impacts in accordance with WA EPA guidelines and impact mitigation hierarchy.

- EPBC Referral Report No. 7 - Commonwealth Matters.

 Section 10.4 assesses potential impacts of the proposal on listed species including sawfish in accordance with the EPBC Act significant impact criteria and impact mitigation hierarchy, and finds no significant or residual impacts in accordance with these criteria.

- EPBC Supplementary Report No. 3 - Commonwealth Environmental Management Plan (C-EMP).

 This includes best practice impact avoidance, mitigation, monitoring and response actions for sawfish in accordance with the impact mitigation hierarchy, as summarized against Element 6 below.

2. Distribution & populations:

Largetooth Sawfish are found globally in all tropical coastal waters and estuarine and river systems in the Atlantic, Indian and Pacific

They have an extremely broad distribution ranging from freshwater bodies up to 400 km inland to coastal and marine waters up to 100 km offshore. They have an ontogenetic shift in habitat

As reported in Section 9.4.5 of <u>EPBC Referral Report No. 2 - Setting & Existing Environment</u>, the upstream areas of the rivers and creeks that discharge into CG provide habitat that is suitable for neonate and juvenile Largetooth Sawfish, and the coastal waters of CG provide habitat that is suitable for adult Largetooth Sawfish.

However, no previously published papers, reports or verifiable data could be found confirming their presence in CG. As outlined above the eDNA sampling and the marine fauna surveys commissioned by BKA found no evidence of their presence in CG.

Key element from the Conservation Advice How addressed by BKA utilisation with neonate and juvenile animals Never-the-less, consistent with the precautionary principle, it is assumed that primarily occurring in the freshwater and less the occasional adult could potentially pass through the POA during saline waters of rivers and estuaries and adult inshore/offshore movements. animals being found in more saline coastal and marine waters. In Australia they are found in coastal waters and estuarine and river systems across the tropical north of the country. The Conservation Advice for this species does not provide a population estimate. 3. Foraging behaviour, diet & critical habitat: There is no overlap of the proposed operation with the upriver habitat that might be used by juvenile and sub-adult sawfish and no potential for direct The Conservation Advice does not contain impacts on iuveniles and sub-adults. information on the foraging behaviour and diet of this species. The DCCEEW SPRAT profile for As outlined above, while there are no confirmed records of this species in CG, this species states that it feeds on fishes and including in the deeper marine waters of the main body of CG where the POA benthic invertebrates. The saw is used to stun is located, consistent with the precautionary principle it is assumed that the schooling fish, such as mullet, and for extracting occasional adult could potentially pass through the POA during molluscs and small crustaceans from the benthic inshore/offshore movements. sediment. Potential impacts on adult sawfish moving through the POA, and proposed The Conservation Advice states that the generally impact prevention, mitigation and monitoring measures are summarised accepted model of movement of Largetooth against Element 5 below. Sawfish is that young are born at the mouths of rivers and then migrate upriver where they spend the first several years of life. As they reach maturity they move out of the rivers and into the marine environment. The Conservation Advice states that the distribution of this species is not known to overlap with any EPBC Act-listed threatened ecological communities. 4. Significance to First Nations people: Please refer: - EPBC Referral Report No. 3 - Traditional Owner Matters. The Conservation Advice states that: EPBC Referral Report No. 6 - Consultation. Fishing of sawfish is a part of traditional fishing practices and historically makes up an BKA has undertaken and continues to undertake comprehensive consultations important part of the diet of coastal indigenous and engagement with the two First Nations groups in the CG area, Balanggarra and Miriuwang-Gajerrong. Neither group has expressed concerns about sawfish species and both groups have issued letters of support for the Indigenous Australians are allowed to take and eat sawfish for personal, domestic or nonproposal, as presented in Annexes to EPBC Referral Report No. 3 - Traditional commercial communal needs Owner Matters. 5. Main threats & potential impacts of the The proposed operation will not contribute to any of the threats listed in the proposed operation: Conservation Advice. It does not include any form of fishing, it will not discharge debris into the marine environment (see section 3.5 below), and it The Conservation Advice lists the main threats to will not degrade or modify the species' habitat. sawfish as: As outlined above the occasional adult could potentially pass through the POA - commercial fishing (especially gill netting and during inshore/offshore movements. As this is an epibenthic species (it swims prawn trawling), near the seabed) there would be some potential of being entrained in the recreational fishing, SPV's drag-head when it is operating. - indigenous fishing, illegal, unreported and unregulated (IUU) The potential for this species to be entrained in the SPV's drag-head is very low for the following reasons: - entanglement in marine debris; and Apparent absence of this species in the POA (based on surveys and - habitat degradation and modification. eDNA sampling). Very low presence of the SPV in CG (1-2 days every 2 weeks with zero presence in CG for 86% of the time during the project lifespan). Very low vessel speed: The SPV will operate at very low speeds (<2 knots) when loading sand in CG, allowing fauna to move away; and improving the effectiveness of MFOA measures. Single drag-head: The SPV will only have a single suction-arm and draghead (standard TSHD's usually have a pair of suction-arms and drag Marine fauna excluder ('turtle tickler chains'): Will be fitted in front of the drag-head.

Key element from the Conservation Advice	How addressed by BKA
6. Research Priorities & Priority Actions: The Conservation Advice lists a number of research priorities and priority actions, which are strategic in nature for adoption by relevant parties involved in the conservation of the species at the national level. The research priorities and priority actions are designed primarily to address the key threats outlined against Element 5 above.	The proposed operation will not contribute to any of the threats listed in the Conservation Advice which are addressed by the research priorities and priority actions.
7. Recovery Plan: The Conservation Advice references a Draft Sawfish and River Sharks Multispecies Recovery Plan, which was published as a final plan in 2015.	Please refer Table 9 in section 3.4 below which presents BKA's response to relevant elements of the Sawfish and River Sharks Multispecies Recovery Plan.

TABLE 7: Conservation Advice for Pristis zijsron (Green Sawfish)

(undated)

Key element from the Conservation Advice How addressed by BKA As an MNES species, BKA has given very high priority to assessing potential impacts of the proposal and developing relevant impact avoidance, mitigation and monitoring measures for this species, as presented for this Element in Table 6 for Largetooth Sawfish above, which is not repeated here for reasons of economy. Image credit: R Pion Adult size: Up to 5 m. 1. Conservation Status: Green Sawfish are listed as Vulnerable under the EPBC Act, making the species a Matter of National Environmental Significance (MNES) under that Act 2. Distribution & populations: As reported in Section 9.4.5 of EPBC Referral Report No. 2 - Setting & Existing Environment, the upstream areas of the rivers and creeks that discharge into CG provide habitat that is suitable for neonate and juvenile The Conservation Advice states that the species Green Sawfish, and the coastal waters of CG provide habitat that is suitable is currently known to be present across northern Australia from Broome in WA to Cairns in for adult Green Sawfish. Queensland where it inhabits less saline riverine However, no previously published papers, reports or verifiable data could be and estuarine and also marine waters, but does not move into purely freshwater areas. found confirming their presence in CG. As outlined above the eDNA sampling and the marine fauna surveys commissioned by BKA found no evidence of The Conservation Advice does not provide a their presence in CG. population estimate. Never-the-less, consistent with the precautionary principle, it is assumed that the occasional adult could potentially pass through the POA during inshore/offshore movements. As per this Element in Table 6 for Largetooth Sawfish above, which is not 3. Foraging behaviour, diet & critical habitat: repeated here for reasons of economy. The Conservation Advice does not contain information on the foraging behaviour and diet of this species. Like all sawfish it feeds on fishes and benthic invertebrates. The saw is used to stun schooling fish, such as mullet, and for extracting molluscs and small crustaceans from the benthic sediment. Like other sawfish, less saline riverine and estuarine areas are used by neonates, juveniles and sub-adults, which migrate to coastal and offshore areas as adults. 4. Significance to First Nations people: As per this Element in Table 6 for Largetooth Sawfish above, which is not repeated here for reasons of economy. The Conservation Advice does not state anything on this element. As per this Element in Table 6 for Largetooth Sawfish above, which is not 5. Main threats & potential impacts of the proposed operation: repeated here for reasons of economy. The Conservation Advice lists the main threats to Green Sawfish as: incidental capture as bycatch in gillnet and trawl fisheries. illegal capture for fins and rostra, habitat degradation through coastal development

Key element from the Conservation Advice	How addressed by BKA
6. Priority Actions: The Conservation Advice lists a number of priority actions, which are strategic in nature for adoption by relevant parties involved in the conservation of the species at the national level. The priority actions are designed primarily to address the key threats outlined against Element 5 above.	The proposed operation will not contribute to any of the threats listed in the Conservation Advice which are addressed by the priority actions.
7. Recovery Plan: The Conservation Advice does not reference a Recovery Plan.	Please refer Table 9 in section 3.4 below which presents BKA's response to relevant elements of the Sawfish and River Sharks Multispecies Recovery Plan.

 TABLE 8: Conservation Advice for <u>Pristis clavata</u> (Dwarf Sawfish)

In effect under EPBC Act from 7 October 2009.

Key element from the Conservation Advice	How addressed by BKA
Image credit: R Kuiter Adult size: Up to 3.2 m. 1. Conservation Status:	As an MNES species, BKA has given very high priority to assessing potential impacts of the proposal and developing relevant impact avoidance, mitigation and monitoring measures for this species, as presented for this Element in Table 6 for Largetooth Sawfish above, which is not repeated here for reasons of economy.
Dwarf Sawfish are listed as <u>Vulnerable</u> under the EPBC Act, making the species a Matter of National Environmental Significance (MNES) under that Act.	
2. Distribution & populations: The Conservation Advice states that the species is currently known to be present across northern Australia from the Pilbara in WA to Cairns in Queensland where it inhabits less saline riverine and estuarine and also marine waters, but does not move into purely freshwater areas. The Conservation Advice does not provide a population estimate.	As reported in Section 9.4.5 of EPBC Referral Report No. 2 - Setting & Existing Environment, the upstream areas of the rivers and creeks that discharge into CG provide habitat that is suitable for neonate and juvenile Dwarf Sawfish, and the coastal waters of CG provide habitat that is suitable for adult Dwarf Sawfish. However, no previously published papers, reports or verifiable data could be found confirming their presence in CG. As outlined above the eDNA sampling and the marine fauna surveys commissioned by BKA found no evidence of their presence in CG. Never-the-less, consistent with the precautionary principle, it is assumed that the occasional adult could potentially pass through the POA during inshore/offshore movements.
3. Foraging behaviour, diet & critical habitat: The Conservation Advice does not contain information on the foraging behaviour and diet of this species. Like all sawfish it feeds on fishes and benthic invertebrates. The saw is used to stun schooling fish, such as mullet, and for extracting molluscs and small crustaceans from the benthic sediment. Like other sawfish, less saline riverine and estuarine areas are used by neonates, juveniles and sub-adults, which migrate to coastal and offshore areas as adults.	As per this Element in Table 6 for Largetooth Sawfish above, which is not repeated here for reasons of economy.
Significance to First Nations people: The Conservation Advice does not state anything on this element.	As per this Element in Table 6 for Largetooth Sawfish above, which is not repeated here for reasons of economy.
5. Main threats & potential impacts of the proposed operation: The Conservation Advice lists the main threats to Dwarf Sawfish as: incidental capture as bycatch in gillnet fishing, IUU fishing, habitat degradation through coastal development.	As per this Element in Table 6 for Largetooth Sawfish above, which is not repeated here for reasons of economy.
6. Priority Actions:	The proposed operation will not contribute to any of the threats listed in the Conservation Advice which are addressed by the priority actions.

Key element from the Conservation Advice	How addressed by BKA
The Conservation Advice lists a number of priority actions, which are strategic in nature for adoption by relevant parties involved in the conservation of the species at the national level. The priority actions are designed primarily to address the key threats outlined against Element 5 above.	
7. Recovery Plan: The Conservation Advice does not reference a Recovery Plan.	Please refer Table 9 in section 3.4 below which presents BKA's response to relevant elements of the Sawfish and River Sharks Multispecies Recovery Plan.

3.4 Species Recovery Plans

- 1. As outlined in section 1.2 above, item 1.3 of DCCEEW's RFI states 'Please demonstrate that the proposal has had regard to relevant conservation advices, and <u>is not inconsistent with recovery plans</u> and/or threat abatement plans, etc.'
- 2. The Australian Government Minister for the Environment may adopt and implement recovery plans for threatened fauna, flora (other than conservation dependent species) and ecological communities listed under the EPBC Act.
- 3. Recovery plans set out research and management actions to stop the decline of, and support the recovery of, listed threatened species or threatened ecological communities. The aim of a recovery plan is to maximise the long-term survival in the wild of a threatened species or ecological community. They should also state how to manage and reduce threatening processes.
- 4. Recovery plans provide a planned and logical framework for responsible government agencies and key interest groups. This helps them to coordinate their work to improve outcomes for threatened species and ecological communities.
- Currently (August 2025) there are two approved species recovery plans, both covering multiple species, that are relevant to key threatened species in the CG area, as follows (web links):
 - Sawfish & River Sharks Multi-species Recovery Plan. 2015. Covers the following species:
 - Speartooth Shark (<u>Glyphis glyphis</u>).
 - Northern River Shark (<u>Glyphis garricki</u>).
 - Largetooth Sawfish (Pristis pristis).
 - Green Sawfish (Pristis zijsron).
 - Dwarf Sawfish (Pristis clavata).
 - Recovery Plan for Marine Turtles in Australia, 2017-2027. Covers the following species:
 - Flatback Turtle (Natator depressus) (the most relevant species to the CG area).
 - Hawksbill Turtle (Eretmochelys imbricata).
 - Green Turtle (Chelonia mydas).
 - Leatherback Turtle (Dermochelys coriacea).
 - Loggerhead Turtle (Caretta caretta).
 - Olive Ridley Turtle (Lepidochelys olivacea).
- 6. Tables 9 and 10 present key elements of each of these two recovery plans respectively, and how these elements have been addressed by BKA for the CG marine sand proposal.

TABLE 9: Sawfish & River Sharks Multi-species Recovery Plan

Published 2015.

NOTE: The Recovery Plan repeats much of the information that is presented in the Conservation Advice for each species that are assessed in section 3.3 above. This is not repeated in this section for reasons of economy.

Key element from the Recovery Plan	How addressed by BKA
1. Species covered: The Recovery Plan covers the following species: Speartooth Shark (Glyphis glyphis). Northern River Shark (Glyphis garricki). Largetooth Sawfish (Pristis pristis). Green Sawfish (Pristis zijsron). Dwarf Sawfish (Pristis clavata). These species have been grouped together in a single recovery plan because of similarity in habitat use, distribution and threats to recovery. All of these species predominantly inhabit the rivers, estuaries and inshore marine habitats of northern Australia.	BKAs' approach to the protection of each of these species is presented in the same order as listed, in Tables 4 to 8 in section 3.3 above, in relation to relevant aspects of the Conservation Advice for each species.
2. Threats to the species: The Recovery Plan is consistent with the Conservation Advice for each species reviewed in section 3.3 above, in that it states that the principal threats to the sawfish and river shark species come from commercial, recreational and indigenous and IUU fishing, habitat degradation and modification, as well as the collection of animals for display in public aquaria and marine debris.	As outlined in Tables 4 to 8 in section 3.3 above the proposed operation will not contribute to any of the threats listed in the Recovery Plan. It does not include any form of fishing, it will not discharge debris into the marine environment, and it will not degrade or modify the species' habitat.
3. Overarching objective of the Recovery Plan: The overarching objective of the Recovery Plan is to assist the recovery of these species in the wild throughout their range in Australian waters by increasing their total population size, with a view to: — improving the population status leading to the removal of these species from the protected species list of the EPBC Act; and — ensuring that anthropogenic activities do not hinder recovery in the near future, or impact on the conservation status of the species in the future.	As outlined in Tables 4 to 8 in section 3.3 above the proposed operation will not contribute to any of the threats listed in the Recovery Plan, and does not pose a risk of significant impact on these species that might affect their population or conservation status, consistent with the EPBC Act significant impact guidelines and criteria. The measures to be implemented by BKA as summarized in section 3.3 above, including the data that will be generated by the proposed monitoring program, will contribute to improved understanding of the species and to their conservation and recovery.
4. Specific objectives of the Recovery Plan: The Recovery Plan lists 10 specific objectives:	
Objective 1: Reduce and, where possible, eliminate adverse impacts of commercial fishing on sawfish and river shark species.	The proposed operation does not involve commercial fishing.
 Objective 2: Reduce and, where possible, eliminate adverse impacts of <u>recreational</u> <u>fishing</u> on sawfish and river shark species. 	The proposed operation does not involve recreational fishing.
Objective 3: Reduce and, where possible, eliminate adverse impacts of indigenous fishing on sawfish and river shark species.	The proposed operation does not involve indigenous fishing.

Key element from the Recovery Plan	How addressed by BKA
Objective 4: Reduce and, where possible, eliminate the impact of IUU fishing on sawfish and river shark species.	The proposed operation does not involve IUU fishing.
Objective 5: Reduce and, where possible, eliminate adverse impacts of <u>habitat</u> degradation and modification on sawfish and river shark species.	The proposed operation will not degrade or modify the species' habitat. The POA is located in deeper, open waters in the central part of CG, away from coastal foraging areas. The sandy seabed within the POA, which is highly dynamic with constantly mobile sand-waves driven by strong tidal currents, does not provide suitable foraging habitat.
Objective 6: Reduce and, where possible, eliminate any adverse impacts of marine debris on sawfish and river shark species noting the linkages with the Threat Abatement Plan for the Impact of Marine Debris on Vertebrate Marine Life.	The proposed operation will not discharge debris into the marine environment (see also section 3.5 below).
Objective 7: Reduce and, where possible, eliminate any adverse impacts of collection for public aquaria on sawfish and river shark species.	The proposed operation does not involve collecting for public aquaria.
Objective 8: Improve the information base to allow the development of a quantitative framework to assess the recovery of, and inform management options for, sawfish and river shark species.	This is not applicable to the proposed operation as it will not contribute to any of the key threats to the species as identified in the Recovery Plan, which are aligned with the 10 objectives.
Objective 9: Develop research programs to assist conservation of sawfish and river shark species.	This is not applicable to the proposed operation as it will not contribute to any of the key threats to the species as identified in the Recovery Plan, which are aligned with the 10 objectives.
Objective 10: Improve community understanding and awareness in relation to sawfish and river shark conservation and management.	This is not applicable to the proposed operation as it will not contribute to any of the key threats to the species as identified in the Recovery Plan, which are aligned with the 10 objectives.
5. Actions to achieve the 10 Objectives: The Recovery Plan outlines recommended actions to achieve the 10 objectives, with associated performance criteria and identification of responsible agencies and potential partners.	The recommended actions are not applicable to the proposed operation as they are aligned with the 10 objectives, which are not applicable to the proposed operation as outlined above. Depending on the recommended action, the responsible agencies are identified as Commonwealth, State and Territory and local government agencies, the commercial and recreational fishing sectors, research institutions, NGOs and First Nations groups, and potential partners also include the commercial and recreational fishing sectors, research institutions, NGOs and First Nations groups.

 TABLE 10: Recovery Plan for Marine Turtles in Australia, 2017-2027

Key element from the Recovery Plan	How addressed by BKA
1. Species covered: The Recovery Plan covers all six species of marine turtle that are found in Australian waters, as follows: Flatback Turtle (Natator depressus) (the most relevant species to the CG area). Hawksbill Turtle (Eretmochelys imbricata). Green Turtle (Chelonia mydas). Leatherback Turtle (Dermochelys coriacea). Loggerhead Turtle (Caretta caretta). Olive Ridley Turtle (Lepidochelys olivacea).	BKA has given priority focus to Flatback Turtles as the most significant species found in the general CG area and given the significant Flatback nesting beach at Cape Domett to the east and just outside of CG. However, the impact prevention, mitigation, monitoring and response measures proposed by BKA apply equally to any species of marine turtle. A detailed description of marine turtles, including Flatback Turtles in the CG area, is presented in section 9.4 of EPBC Referral Report No. 2 - Setting & Existing Environment , supported by Annex 12 to that report - Domett Turtle Data Report . Potential impacts of the proposed operation on Flatback Turtles are assessed in Section 10.3.2 of EPBC Referral Report No. 4 - Impact Assessments in accordance with WA EPA guidelines and impact mitigation hierarchy, and find no significant or residual impacts in accordance with those guidelines. Potential impacts of the proposed operation on Flatback Turtles are assessed in Section 10.2 of EPBC Referral Report No. 7 - Commonwealth Matters in accordance with EPBC Act significant impact criteria and impact mitigation hierarchy, and finds no significant or residual impacts in accordance with those criteria. Additional information on marine turtle issues is presented in Section 2 - Current Speeds in the POA & Turtle Swimming Speeds, and Section 3 - Cape Domett , of <a (c-emp)"="" -="" 3="" commonwealth="" environmental="" epbc="" href="EPBC Referral Supplementary Report No. 4 - Additional Information. Proposed impact prevention, mitigation, monitoring and response measures for marine turtles are presented in EPBC Supplementary Report No. 3 - Commonwealth
2. Threats to the species: The Recovery Plan applies a risk assessment approach and states that the risk posed by anthropogenic threats to marine turtle stocks varies depending on the habitats they occupy, timing of habitat occupancy, life cycle stage affected, abundance and trends in nesting and foraging numbers, and the management and mitigation currently in place. The Recovery Plan identifies the following main anthropogenic threats to marine turtles:	
- climate change and variability;	The proposed operation will not contribute to climate change and variability to a degree that could affect marine turtles. Section 12 of EPBC Referral Report No. 5 - Impact Assessments assesses greenhouse gas (GHG) emissions from the proposed operation in accordance with WA EPA guidelines, and finds that emissions will not exceed the WA EPA trigger level of 100,000 tonnes CO2-e in any year, and therefore does not trigger assessment under the WA Environmental Protection Act. The SPV will comply with Annex VI (Air Pollution) of the MARPOL Convention and the implementing Australian regulations (AMSA Marine Order 97). These regulations set strict standards and limits on GHG emissions from ships, and require ships to implement a range of on-board energy efficiency and emissions reduction strategies and plans, including having an IMO-compliant ship-specific Energy Efficiency Design Index (EEDI) and Shipboard Energy Efficiency Management Plan (SEEMP).
marine debris;	The proposed operation will not discharge debris into the marine environment (see also section 3.5 below).
chemical and terrestrial discharge;	The proposed operation will not contribute to chemical and terrestrial discharge.

Key element from the Recovery Plan	How addressed by BKA
international take;	The proposed operation will not contribute to international take.
terrestrial predation;	The proposed operation will not contribute to terrestrial predation.
- fisheries bycatch;	The proposed operation will not contribute to fisheries bycatch.
light pollution;	Potential impacts of light emissions from the SPV on nesting and hatching turtles in the CG area are assessed in EPBC Supplementary Report No. 1 - Light Assessment in accordance with both the National Light Pollution Guidelines for Wildlife (DCCEEW 2023) and WA EPA requirements, and finds no significant impacts.
	The SPV will be permanently fitted with turtle safe lighting as specified in DCCEEW (2023) and other light impact prevention, mitigation and monitoring measures will be implemented as outlined for CEO 10 - SPV Lighting in EPBC Supplementary Report No. 3 - Commonwealth Environmental Management Plan (C-EMP).
habitat modification through infrastructure/coastal development and dredging and trawling;	The proposed operation will not cause habitat modification through infrastructure/coastal development as it does not involve any form of infrastructure/coastal development – it is a 100% vessel-based marine operation.
	The proposed operation will not involve trawling.
	The proposed sand-sourcing will involve a form of dredging, with the SPV being based on the design principles of a Trailer Suction Hopper Dredger (TSHD), albeit with only a single suction-arm and drag-head (standard TSHD's usually have a pair of suction-arms and drag-heads).
	The proposed sand sourcing will not modify habitat that is significant to marine turtles. Sand sourcing will be restricted to the POA which is located in the deeper open waters of the main body of CG, with an average depth of -20.6 m LAT and a max depth of -44 m LAT. The seabed in the POA comprises highly dynamic, constantly mobile sand-waves driven by extremely strong tidal currents, and does not provide suitable foraging or inter-nesting habitat for marine turtles. A detailed description of the environmental conditions and lack of benthic biota in the POA is provided in sections 5, 6.4.2, 6.4.3 and 6.4.4 of 6.4.4.8 of EPBC Referral Report No. 2 - Setting & Existing Environment.
	Although the environmental conditions within CG do not provide suitable internesting habitat, DCCEEW has declared a Flatback Turtle inter-nesting 'buffer' Biologically Important Area (BIA) for a 60 km radius around Cape Domett and Lacrosse Island, which includes the waters within CG. This is discussed in detail in the response to Element 7 below.
	The most important turtle habitat in the CG area is a major Flatback Turtle nesting beach at the Cape Domett Seaward Beach, located outside and to the east of CG, and lesser nesting beaches at Turtle Bay on Lacrosse Island, at Turtle Bay West on the seaward coast outside of CG, west of Cape Dussejour, and a nesting site located on stranded sand ridges behind mangroves at Barnett Point, inside CG. Peak nesting is in August-September, in contrast to Flatback Turtles on the west coast of WA, where peak nesting is in Nov-Jan (Whiting et al 2008). A detailed description of the turtle nesting beaches in the CG area, including the results of aerial drone surveys, is presented in section 9.2.5 of EPBC Referral Report No. 2 - Setting & Existing Environment, supported by Annex 12 to that report - DBCA Cape Domett Turtle Data Report.
	Potential impacts of the proposed operation on the turtle nesting beaches in the CG area are assessed in detail in Section 5 - Sediment Transport & Beach Processes, of EPBC Referral Report No. 8 - Full Modeling Report. This included high-resolution aerial drone LiDAR surveys of the beaches, analysis of historical beach dynamics using satellite imagery, characterisation of sediment supply processes at each beach, and numerical modelling to predict potential changes at the beaches from potential changes to sediment supply from the proposed sand sourcing within CG. The assessment found that the proposed operation will not affect beach processes. Never-the-less, as a precaution, should the proposed operation go ahead, monitoring of the beaches is proposed, including regular high-resolution aerial drone LiDAR surveys, as outlined in CEO 9: Coastal Processes & Beaches of EPBC

Key element from the Recovery Plan	How addressed by BKA
	Supplementary Report No. 3 - Commonwealth Environmental Management
	<u>Plan (C-EMP).</u>
– indigenous take;	The proposed operation will not contribute to indigenous take.
 vessel disturbance; 	The two main potential vessel disturbance impacts of the proposed operation on marine turtles are: - Vessel strike: Potential vessel strike from the Sand Production Vessel (SPV) on a turtle swimming at or near the sea surface; and - Drag-head entrainment: Potential entrainment in the SPV's draghead, in the unlikely event that a marine turtle is sitting on the seabed in the path of the drag-head when it is operating. Potential impacts are assessed in Section 10.3.2 - Flatback Turtles of EPBC Referral Report No. 4 - Impact Assessments in accordance with WA EPA guidelines and impact mitigation hierarchy, and in Section 10.2 - Specific Assessment for Flatback Turtles of EPBC Referral Report No. 7 - Commonwealth Matters in accordance with the EPBC Act significant impact
	criteria and impact mitigation hierarchy. The assessments find no significant or residual impacts in accordance with respective guidelines, criteria and the impact mitigation hierarchy, including through the application of the following impact prevention, mitigation, monitoring and response measures.
	Vessel strike prevention, mitigation, monitoring & response measures:
	Potential vessel strikes will be prevented, mitigated and monitored as outlined in CEO 7 - Vessel Strikes in EPBC Supplementary Report No. 3 - Commonwealth Environmental Management Plan (C-EMP), as follows:
	<u>CEO 7: Vessel Strikes</u> : No significant negative impacts are caused to populations of surface-dwelling marine fauna in CG from vessel strikes by the SPV.
	Impact prevention: Very low likelihood of encounters due to: Very low occurrence of these species in the POA (as indicated by lack of suitable seabed habitat, extreme tidal currents, dedicated site survey results and analysis of satellite tracking data). Very low presence of the SPV in CG (1-2 days every 2 weeks with zero presence in CG for 86% of the time during the project lifespan). SPV Marine Fauna Observation & Avoidance (MFOA) measures (with TOs): Very low vessel speed: The SPV will operate at very low speeds (<2 knots) when loading sand in CG, allowing fauna to move away; and improving the effectiveness of MFOA measures. Western entry/exit route: Restricting entry & exit of the SPV into and out of CG to West Entrance, furthest from the main nesting beach at Cape Domett.
	Impact mitigation: - SPV MFOA measures (with TOs) (this is both an impact prevention & mitigation measure). - Very low speed: The SPV will operate at very low speeds (<2 knots) when loading sand in CG, allowing fauna to move away; and improving the effectiveness of MFOA measures (this is both an impact prevention & mitigation measure). Trigger Criteria (TCs), Trigger Response Actions (TRAs), Threshold Criteria (THCs), Threshold Contingency Actions (TCAs) and monitoring and
	reporting measures for CEO 7 are specified in the Draft EMP, in accordance with WA EPA criteria, which DCCEEW advised is the accepted template for this proposal. Drag-head entrainment prevention, mitigation, monitoring & response
	measures: Potential drag-head entrainment will be prevented, mitigated and monitored as outlined in CEO 11 - Drag-head Entrainment in EPBC Supplementary Report No. 3 - Commonwealth Environmental Management Plan (C-EMP) as follows:

Key element from the Recovery Plan	How addressed by BKA
	CEO 11: Drag-head Entrainment: No significant negative impacts are caused to populations of large epibenthic species in CG from entrainment in the SPV's drag-head.
	Impact prevention: Very low likelihood of encounters due to: Very low occurrence of marine turtles on the seabed in the POA (as indicated by lack of suitable seabed habitat, extreme tidal currents, dedicated site surveys results and analysis of satellite tracking data). Very low presence of the SPV in CG (1-2 days every 2 weeks with zero presence in CG for 86% of the time during the project lifespan). Single drag-head: The SPV will only have a single suction-arm and drag-head (standard TSHD's usually have a pair of suction-arms and drag-heads). SPV MFOA measures (with TOs). Very low speed: The SPV will operate at very low speeds (<2 knots) when loading sand in CG, allowing fauna to move away; and improving the effectiveness of MFOA measures.
	Impact mitigation: - Soft-start procedure: This involves slowly lowering the draghead to the seabed and starting at low pump revolutions, providing opportunity for any marine fauna on the seabed to move away. This is a recognized mitigation measure in the Recovery Plan and is has been accepted as best practice in dredging projects across marine turtle areas of Australia for over ten years Marine fauna excluder ('turtle tickler chains'): Will be fitted in front of the drag-head. This is a recognized mitigation measure in the Recovery Plan and is has been accepted as best practice in dredging projects across marine turtle areas of Australia for over ten years SPV MFOA measures (with TOs) (this is both an impact prevention & mitigation measure) Very low speed: The SPV will operate at very low speeds (<2 knots) when loading sand in CG, allowing fauna to move away; and improving the effectiveness of MFOA measures (this is both an impact prevention & mitigation measure).
	Trigger Criteria (TCs), Trigger Response Actions (TRAs), Threshold Criteria (THCs), Threshold Contingency Actions (TCAs) and monitoring and reporting measures for CEO 11 are specified in the Draft EMP, in accordance with WA EPA criteria, which DCCEEW advised is the accepted template for this proposal.
	Enhanced measures during peak-nesting season:
	While the above measures are assessed as being more than adequate for preventing and mitigating the potential for significant impacts in all seasons, as an additional precaution enhanced vessel strike and drag-head prevention and mitigation measures are proposed during the peak Flatback Turtle nesting season in the CG area (Aug-Sept), including: - Very low presence of the SPV in CG during this period (4 loading cycles of up to 2 days each = max of 8 days presence during the season). - Spatial restriction on sand-sourcing operations to the western half of the POA (furthest from the main nesting beach at Cape Domett) during the season (please refer map of the proposed restricted area in EPBC Supplementary Report No. 3 - Commonwealth Environmental Management Plan (C-EMP). - Doubling the MFOA effort during the season (from two active observers to four active observers) (pls refer the C-EMP for details of the proposed MFOA measures).
noise interference;	Underwater Noise:
	The design and construction of the SPV will include relevant best-practice noise reduction measures in accordance with the <i>IMO Underwater Radiated Noise Guidelines</i> (IMO 2023), which constitute the international standard.
	Potential impacts of underwater noise are assessed in <u>EPBC Supplementary</u> Report No. 2 - Noise Assessment. This includes detailed modelling of predicated noise emissions from the SPV and assessment of potential auditory

Key element from the Recovery Plan	How addressed by BKA
	injury and behavioural impacts on marine turtles, in accordance with the US NMFWS criteria (as required by WA EPA), using a risk assessment approach, and finds that potential impacts are negligible. Despite the assessment that potential impacts are negligible, BKA proposes to undertake monitoring of underwater noise in CG during commencement of operations to assess compliance with the findings of the Noise Assessment, as described in EPBC Supplementary Report No. 3 - Commonwealth Environmental Management Plan (C-EMP).
recreational activities; and	The proposed operation will not contribute to impacts from recreational activities.
 disease and pathogens. 	The proposed operation will not contribute to significant impacts from disease and pathogens in marine turtles (pls refer section 3.2 above).
3. Long-term recovery objective: The long-term recovery objective of the Recovery Plan is to minimise anthropogenic threats to allow for the conservation status of marine turtles to improve so that they can be removed from the EPBC Act threatened species list.	BKA is already contributing to the long-term recovery objective as reported in Annex 12 - <u>DBCA Cape Domett Turtle Data Report</u> of <u>EPBC Referral Report No. 2 - Setting & Existing Environment</u> , including: - Entering into a data-sharing agreement with the WA Department of Biodiversity Conservation & Attractions (DBCA) and undertaking analysis and reporting of ten-years of data from DBCA's long-term monitoring program at the Cape Domett Seaward Beach, to help inform the improved
	management of marine turtles. - Undertaking dry- and wet-season aerial drone surveys of all potential turtle nesting sites in the CG area, and expanding the identification and characterization of nesting sites in the area.
	 Undertaking dry- and wet-season marine fauna surveys, including for marine turtles, throughout CG area, covering over 820 km of transects in each survey, to provide data to help inform the improved management of marine turtles.
	Should the proposal proceed, BKA would seek to further support the long-term recovery objective, in cooperation with relevant agencies and local First Nation's peoples, including:
	 Supporting the expansion of DBCA's current long-term monitoring prograr at the Cape Domett Seaward Beach to other nesting sites in the CG area.
	 Supporting satellite tagging and movement tracking of marine turtles in the CG area.
	 Implementing proposed impact prevention, mitigation, monitoring and response measures for marine turtles outlined in <u>EPBC Supplementary</u> <u>Report No. 3 - Commonwealth Environmental Management Plan (C-EMP)</u>
	 Making all data from the proposed Marine Fauna Observation & Avoidance (MFOA) program publicly available (e.g. on a web site), to further inform the improved management of marine turtles.
4. Interim recovery objectives:	
The plan suggests that the long-term recovery objective is unlikely to be achieved during the ten-year life of the plan (2017-2027), and therefore sets four interim objectives and associated actions for the life of the plan (to 2027), as follows:	
 Interim Objective 1: Current levels of legal and management protection for marine turtles are maintained or improved both domestically and throughout the migratory range of Australia's marine turtles. 	This Interim Objective is primarily the responsibility of government agencies and research bodies.
	-

Key element from the Recovery Plan	How addressed by BKA
 Interim Objective 2: The management of marine turtles is supported. 	BKA is already contributing to improving the management of marine turtles, as outlined in the response to the Element 3 - Long-term recovering objective, above. Should the proposal proceed, BKA would seek to further support improving the management of marine turtles, in cooperation with relevant agencies and local First Nation's peoples, as outlined in the response to the Element 3 - Long-term recovering objective, above.
 Interim Objective 3: Anthropogenic threats are demonstrably minimised. 	The proposed impact prevention, mitigation, monitoring and response measures for marine turtles outlined in <u>EPBC Supplementary Report No. 3 - Commonwealth Environmental Management Plan (C-EMP)</u> and as summarized throughout this table, will prevent and minimize anthropogenic threats to marine turtles in the CG area, as far as they relate to the proposed operation.
 Interim Objective 4: Trends at index beaches, and population demographics at important foraging grounds are described. 	BKA is already contributing to improving the understanding of Flatback Turtle nesting and population trends at the Cape Domett Seaward Beach and other beaches in the CG area as, as outlined in the response to Element 3 - Long-term recovering objective, above. Should the proposal proceed, BKA would seek to further support understanding of nesting and population trends as also outlined in the response to Element 3 - Long-term recovering objective, above.
6. Actions: The Recovery Plan includes a number of Actions that are designed to address the main threats to marine turtles as listed against Element 2 above and support achievement of the Interim Objectives as listed against Element 5 above, divided into A. Assessing & addressing threats and B. Enabling & measuring recovery, as follows:	NOTE: There is repetition in the responses to Element 6 below as the Actions largely mirror the Interim Objectives in Element 5 above. However, each Action is included and addressed in turn so as to prove a complete picture of how BKA has addressed / will address all aspects of the Recovery Plan.
A Actions - Assessing & addressing threats:	
A1. Maintain and improve efficacy of legal and management protection.	This Action is primarily the responsibility of government agencies and research bodies.
 A2. Adaptively manage turtle stocks to reduce risk and build resilience to climate change and variability. 	As per response against climate change in Element 2 above, the proposed operation will not contribute to climate change and variability to a degree that could affect marine turtles.
A3. Reduce the impacts from marine debris.	As per response against marine debris in Element 2 above, the proposed operation will not discharge debris into the marine environment (see also section 3.5 below).
 A4. Minimise chemical and terrestrial discharge: 	As per response against chemical and terrestrial discharge in Element 2 above, the proposed operation will not contribute to chemical and terrestrial discharge.
 A5. Address international take within and outside Australia's jurisdiction: 	As per response against international take in Element 2 above, the proposed operation will not contribute to international take.
A6. Reduce impacts from terrestrial predation:	As per response against terrestrial predation in Element 2 above, the proposed operation will not contribute to terrestrial predation.
 A7. Reduce international and domestic fisheries bycatch: 	As per response against fisheries bycatch in Element 2 above, the proposed operation will not contribute to fisheries bycatch.

Key element from the Recovery Plan	How addressed by BKA
A8. Minimise light pollution:	As per response against light pollution in Element 2 above, the proposed operation will not cause impacts on marine turtles from light pollution.
A9. Address the impacts of coastal development/infrastructure and dredging and trawling:	As per response against this item in Element 2 above, the proposed operation will not cause habitat modification through infrastructure/coastal development and will not involve trawling. As per response against this item in Element 2 above, the proposed sand-sourcing will involve a form of dredging within the POA only, and this will not cause impacts on habitat that is significant for marine turtles.
A10. Maintain and improve sustainable Indigenous management of marine turtles:	As per response against indigenous take in Element 2 above, the proposed operation will not contribute to indigenous take. Should the proposal proceed, BKA would seek to further support improving the management of marine turtles, in cooperation with relevant agencies and local First Nation's peoples, as outlined in the response to the Element 3 - Long-term recovering objective, above.
B Actions - Enabling & measuring recovery:	
B1. Determine trends at index beaches:	As per response against Interim Objective 4 under Element 5 above, BKA is already contributing to improving the understanding of Flatback Turtle nesting and population trends at the Cape Domett Seaward Beach and other beaches in the CG area. Should the proposal proceed, BKA would seek to further support
	understanding of nesting and population trends as also outlined in the response to Element 3 - Long-term recovering objective, above.
B2. Understand population demographics at key foraging grounds:	CG is not a foraging ground for marine turtles so this Action is not directly relevant.
B3. Address information gaps to better facilitate the recovery of marine turtle stocks:	As also outlined in the response to Element 3 - Long-term recovering objective, and other response above, BKA is already contributing to addressing information gaps to better facilitate the recovery of marine turtle stocks, and will continue to do so, should the proposed operation proceed.
7. Biologically Important Areas (BIA's): The Recovery Plan describes BIAs as areas where protected species display biologically important behaviour, such as breeding, foraging, resting and migration, and identifies various BIAs	An inter-nesting area is where female turtles 'rest' between each egg-laying effort on the nesting beaches, regaining energy and strength for the next egg-laying effort, which are energetically very demanding. Flatback Turtle internesting periods range from a few days up to 13 days, during which time they spend most of their time resting on the seabed, and they do not feed at all during inter-nesting (Whiting et al 2008).
around Australia for all six marine turtle species. Although the environmental conditions within CG do not provide suitable inter-nesting habitat (very strong tidal currents, deeper waters etc), a Flatback Turtle inter-nesting 'buffer' BIA is designated for a 60 km radius around Cape Domett and Lacrosse Island, which includes the waters within CG.	A detailed description of Flatback Turtles and discussion of the BIA in the CG area is presented in section 9.4.2 of <u>EPBC Referral Report No. 2 - Setting & Existing Environment</u> , supported by Annex 12 to that report - <i>DBCA Cape Domett Turtle Data Report</i> .
	Potential impacts of the proposed operation on Flatback Turtles are assessed in Section 10.3.2 of <u>EPBC Referral Report No. 4 - Impact Assessments</u> in accordance with WA EPA guidelines and impact mitigation hierarchy, and find no significant or residual impacts in accordance with those guidelines.
	Potential impacts of the proposed operation on Flatback Turtles are assessed in Section 10.2 of EPBC Referral Report No. 7 - Commonwealth Matters in accordance with EPBC Act significant impact criteria and impact mitigation hierarchy, and finds no significant or residual impacts in accordance with those criteria.
	As outlined in those reports, and particularly in Section 10.2 of EPBC Referral Report No.7 - Commonwealth Matters, the 60 km radius for the Cape Domett inter-nesting BIA is based on satellite tracking from the Pilbara region of WA, located ~ 1,500 km to the west of CG and comprising a different population of Flatback Turtles than those found in the CG area, and which, as outlined above, nest in the opposite season. The tracking showed that Pilbara female Flatbacks can move 'up to' 60 km offshore during inter-nesting, but typically stay much closer (~3 km) to the nesting beach, to conserve energy (Whittock

Key element from the Recovery Plan	How addressed by BKA
	et al 2014). Application of the 60 km radius for the Cape Domett inter-nesting BIA did not consider the characteristics and behaviour of the local population, which are different from the Pilbara, and also did not consider local environmental conditions in CG, which are also different from the Pilbara
	The environmental conditions inside CG, and especially in the POA, including extreme tidal current velocities, deeper waters (up to -44 m LAT with an average depth of -20.6 m) and highly-dynamic, constantly moving sand waves on the seabed, make this area unsuitable for inter-nesting resting by marine turtles.
	Regaining energy during inter-nesting intervals requires no significant net loss of energy reserves as a result of energy expended, including any energy expended from swimming that might be required against currents in the area. Like all marine turtles, Flatback Turtles do not feed during inter-nesting intervals, so energy expenditure must come from stored fat reserves during this period, while also retaining sufficient energy for the following egg laying efforts (which have high energy demand) (Whittock pers. comms., 2025). Therefore, should a turtle be required to expend excess energy during an internesting interval, for example in order to swim against currents in order to remain on the seabed in an area, the 'resting' benefits of inter-nesting would be negated.
	The main nesting beaches in the CG area are located on the seaward coast and face out to sea. After each nesting event Flatbacks would most likely head straight offshore to the inner waters of Joseph Bonaparte Gulf, where current velocities are less and conditions are more favourable than within CG, for their inter-nesting rest, before coming back to the beach again.
	Section 2 - Current Speeds in the POA & Turtle Swimming Speeds of EPBC Referral Supplementary Report No. 4 - Additional Information, presents an analysis of measured current speeds in the POA against typical swim speeds of adult Flatback Turtles. The analysis finds that it is unlikely that Flatback Turtles could effectively rest on the sandy seabed in the POA between nesting attempts, due to the relatively strong near-bed currents. The analysis also finds that based on the spatial distribution of current speeds in the CG area, it is likely that Flatback Turtles would choose an area with lower current speeds for inter-nesting resting (e.g. on the east side of CG, to the north or south of Lacrosse Island and adjacent to / offshore from the nesting beaches), and not in the main body of CG and especially not in the POA. This is borne out by site surveys and tracking of turtles in the CG area.
	Section 3 - Analysis of Turtle Satellite Tracking - Cape Domett, of EPBC Referral Supplementary Report No. 4 - Additional Information, presents an analysis of two previous satellite tagging and tracking programs of nesting female Flatback Turtles at Cape Domett, one in June 209 and one from August 2025 to August 2027, as reported on www.seaturtle.org . A total of 16 turtles were fitted with satellite trackers, comprising and five in the initial and 11 in the later study. The analysis shows that:
	 All 16 tracked turtles undertook inter-nesting movements immediately offshore from Cape Domett until the end of nesting, whereafter they headed further offshore into Joseph Bonaparte Gulf, then either NE towards Darwin and locations in the Arafura Sea beyond, or NW towards the Timor Sea and locations offshore from the West Kimberley.
	Eleven of the 16 tracked turtles do not appear to have entered CG.
	 Two of the 16 tracked turtles appear to have entered CG, but on the far eastern side only, close to the coast near to Cape Domett, and do not appear to have entered the POA.
	 Two of the 16 tracked turtles may have 'possibly' entered CG, although the low resolution of the maps makes this difficult to ascertain, and again on the far eastern side only, close to the coast near to Cape Domett, and they do not appear to have entered the POA.
	Only one of the 16 tracked turtles appears to have crossed the south- eastern corner of the POA.
	The analysis therefore supports the assessment that the waters inside CG and especially in the POA do not provide suitable inter-nesting conditions, that most turtles head straight offshore to the inner waters of Joseph Bonaparte Gulf for their inter-nesting rest, and the few that do enter CG remain close to the coast on the far eastern side, nearest to Cape Domett, where currents are less.

Key element from the Recovery Plan	How addressed by BKA
	The dry- and wet-season marine fauna surveys commissioned by BKA as reported in EPBC Referral Report No. 2 - Setting & Existing Environment - ANNEX 14 - Marine Fauna Surveys Report support this analysis further, with very low numbers of sightings of marine turtles in CG overall, and even less in the POA, as follows:
	 Dry-season survey (late July 2023 - near peak nesting period): Five Flatback Turtle sightings (three near Cape Domett where the main nesting beach is, one near Adolphus Island and one on the west side of CG and none on the POA). Seven unidentified turtle sightings (one near Cape Domett, one near Adolphus Island, one on the west side of CG, one on the east side of CG, two near Lacrosse Island and one within the POA).
	Wet-season survey (February 2024): Two unidentified turtle sightings in CG, one inside the POA, and no other sightings.
	Only one turtle was observed in the POA on each survey, both unidentified species. It should be noted that different sightings could be the same individual(s), so the actual number of turtles may be less than the number of sightings. These are very low numbers of on-water sightings considering the very large area covered (over 820 km per survey), especially in late July 2023 near the peak nesting season, when hundreds of tracks and nests were observed on the nesting beaches.
	These low on-water sighting numbers further indicate that the area within CG may not be significant as an inter-nesting area by Flatback Turtles, despite the 60 km radius of the inter-nesting BIA extending inshore over CG.
	Never-the-less, there is always a possibility of a turtle passing through the POA when the SPV is operating there, presenting the possibility of either a vessel strike if the turtle is on or near the sea surface, or of being entrained in the SPV's drag-head in the unlikely event that the turtle is on the seabed. The proposed impact prevention, mitigation, monitoring and response actions for potential vessel strikes and drag-head entrainment, as well as proposed enhanced measures during peak nesting season, are described against 'vessel disturbance' under Element 2 above, and are not repeated here for reasons of economy.
8. Oil spill risk:	The proposal does not involve any oil and gas installations.
The Recovery Plan states that for the Cape Domett Flatback Turtle stock, potential spills are a concern due to increasing number of oil and gas installations occurring along the Western Australian coast.	There are no oil installations offshore from Cape Domett and the closest offshore gas installation is the Black Tip well located over 100 km offshore from Cape Domett, which produces LNG and not oil and is operated by ENI Australia.
The Recovery Plan identifies a priority action for the recovery of the Cape Domett stock as:	The reference to 'slow to recover habitats, e.g. nesting beaches' is scientifically incorrect. Sand beaches that become oiled are amongst the fastest environments to recover from oiling, and are highly amenable to physical cleaning which can speed up recovery even more quickly.
 Ensure that spill risk strategies and response programs include management for turtles and their habitats, particularly in reference to slow to recover habitats, e.g. nesting beaches and important foraging grounds.' 	BKA has given extremely high priority to preventing a potential oil spill from the SPV when it is operating in CG, and to implementing best practice spill response, containment, clean-up and mitigation and recovery measures, with very high priory placed on the protection of turtle nesting sites in the CG area (which are mostly located outside of CG).
	Potential oil spills are assessed in Annex 2 - Shipping & Oil Spill Risk Assessment of EPBC Referral Report No. 4 - Impact Assessments and finds that risk is low, including through the application of best-practice impact prevention and mitigation measures.
	Potential oil spills will be prevented, mitigated and responded to as described in CEO 2 - Oil Spills in EPBC Supplementary Report No. 3 - Commonwealth Environmental Management Plan (C-EMP).

3.5 Species Threat Abatement Plans

- 1. As outlined in section 2 above, item 1.3 of DCCEEW's RFI states 'Please demonstrate that the proposal has had regard to relevant conservation advices, and is not inconsistent with recovery plans and/or threat abatement plans, etc.'
- 2. Threat abatement plans are developed by DCCEEW and approved by the Australian Government environment minister, to establish a national framework to guide and coordinate Australia's response to key threatening processes for specific threatened species and ecological communities, registered under the EPBC Act. The plans identify research, management and other actions needed to ensure the long-term survival of native species and ecological communities affected by key threatening processes.
- 3. Currently (August 2025) there is only one approved threat abatement plan that relates to coastal and marine species:
 - Threat Abatement Plan for the impacts of marine debris on the vertebrate wildlife of Australia's coasts and oceans (2018) (short title Marine Debris Threat Abatement Plan).
- 4. Under subsection 279(2) of the EPBC Act, the Australian Government environment minister is required to review threat abatement plans at least every five years. However, at August 2025 the 2018 *Marine Debris Threat Abatement Plan* has not yet been reviewed.
- 5. The plan identifies discharges of garbage, including plastics and other debris, from vessels as one of many sources of marine debris that can impact on marine and coastal vertebrate wildlife.
- 6. The plan identifies compliance of vessels with Annex V of the *International Convention for the Prevention of Pollution from Ships* (MARPOL Convention), which regulates garbage pollution from ships, as being the key action for addressing marine debris from vessels.
- 7. The SPV will not discharge garbage or any other forms of debris into the marine environment. It will comply in full with Annex V of the MARPOL Convention. All garbage produced on board the SPV (e.g. from the day-to-day domestic activities of the crew) will be kept on-board the SPV and managed in accordance with a MARPOL-compliant shipboard garbage management plan. All garbage will be discharged to MARPOL-compliant port waste reception facilities at the sand delivery port (Singapore) for appropriate treatment, including recycling where relevant. Details are outlined against CEO 3 Marine Debris, in EPBC Supplementary Report No. 3 Commonwealth Environmental Management Plan (C-EMP).

4. RESPONSE TO RFI ITEM 2 - ECONOMIC & SOCIAL MATTERS

- 1. As outlined in section 2 above, Attachment B to DCCEEW's RFI letter of 16 July 2025 requests BKA to provide information on economic and social matters relating to the proposal.
- 2. An email from BKA's lead environmental consultant to DCCEW on 16 July 2025 pointed out that information on economic and social matters had already been provided, in particular in section 11.3 of Referral Report No 2, sections 8 and 9 of Referral Report No 3 and section 13 of Referral Report No 4, and queried why information that had already been provided would need to be provided again under an RFI.
- 3. In the response from DCCEEW dated 18 July 2025 it was stated:
 - 'Thank you for directing us to the abovementioned sections. The department is satisfied with the information at paragraphs 11-12 of section 13.3 Impact Assessment and section 13.5 Likely Environmental Outcomes of Referral Report No 4, and section 9 of Referral Report No 3. Please simply address that point of the RFI accordingly'
- 4. This item is therefore deemed to have been addressed and that no further action is required by BKA.

5. RESPONSE TO RFI ITEM 3 - ENVIRONMENTAL MANAGEMENT PLAN

- As outlined in section 2 above, Attachment B to DCCEEW's RFI letter of 16 July 2025 provides detailed comments on Draft 1 of the proposed Environmental Management Plan (EMP) for the proposed operation that had been submitted to DCCEEW for review on 10 June 2025.
- Draft 2 of the proposed EMP has been developed, addressing DCCEEW's review comments, and this is submitted separately as <u>EPBC Supplementary Report No. 3 - Commonwealth Environmental Management Plan (C-EMP)</u>, along with a table listing how each of DCCEEW's review comments have been addressed.

REFERENCES

References Cited:

- Brown, A.M., Smith, J., Salgado-Kent, C., Marley, S., Allen, S.J., Thiele, D., Bejder, L., Erbe, C. & Chabanne, D. (2017). Relative
 abundance, population genetic structure and acoustic monitoring of Australian snubfin and humpback dolphins in regions within
 the Kimberley. Report of Project 1.2.4 prepared for the Kimberley Marine Research Program, Western Australian Marine Science
 Institution, Perth, Western Australia, 61pp plus appendices.
- Brown, A.M., Bejder, L., Pollock, K. H. and Allen, S. J. (2016) Site-Specific Assessments of the Abundance of Three Inshore Dolphin Species to Inform Conservation and Management. Front. Mar. Sci. 3:4.doi: 10.3389/fmars.2016.00004
- Commonwealth of Australia (2009). National Assessment Guidelines for Dredging.
- DCCEEW (2024). National Guidelines for the Survey of Cetaceans, Marine Turtles and Dugong.
- DCCEEW (2023). National Light Pollution Guidelines for Wildlife.
- DCCEEW (2021). EPBC Act significant impact guidelines.
- Kyne P.M., et al (2021) Molecular analysis of newly-discovered geographic range of the threatened river shark Glyphis glyphis reveals distinct populations. Report to the National Environmental Science Program, Marine Biodiversity Hub. Charles Darwin University and CSIRO.
- Kyne P.M. (2020). Sampling in 11 rivers yields expanded range and population size estimate for the Northern River Shark. Marine Biodiversity Hub National Environmental Science program: Fact Sheet June 2020.
- Whiting, A., Thomson, A., Chaloupka, M., & Limpus, C. (2008). Seasonality, abundance and breeding biology of one of the largest populations of nesting flatback turtles, Natator depressus: Cape Domett, Western Australia. Australian Journal of Zoology, 2008(5), 297-303.
- Whittock, Paul & Pendoley, Kellie & Hamann, Mark. (2014). Inter-nesting distribution of flatback turtles Natator depressus and industrial development in Western Australia. Endangered Species Research. 26. 25-38. 10.3354/esr00628.

Cited Referral Documents:

- EPBC Referral Report No. 2 Setting & Existing Environment.
- EPBC Referral Report No. 2 Setting & Existing Environment ANNEX 11 Sediment Contamination Assessment.
- EPBC Referral Report No. 2 Setting & Existing Environment ANNEX 12 DBCA Cape Domett Turtle Data Report.
- EPBC Referral Report No. 2 Setting & Existing Environment ANNEX 14 Marine Fauna Surveys Report.
- EPBC Referral Report No. 3 Traditional Owner Matters.
- EPBC Referral Report No. 4 Impact Assessments.
- EPBC Referral Report No. 6 Consultation.
- EPBC Referral Report No. 7 Commonwealth Matters.
- EPBC Referral Report No. 8 Full Modeling Report
- EPBC Supplementary Report No. 1 Light Assessment.
- EPBC Supplementary Report No. 2 Noise Assessment.
- EPBC Supplementary Report No. 3 Commonwealth Environmental Management Plan (C-EMP).
- EPBC Referral Supplementary Report No. 4 Additional Information.

Conservation Advice documents:

- Conservation Advice for Orcaella heinsohni (Australian Snubfin Dolphin), March 2025.
- Conservation Advice for Sousa sahulensis (Australian Humpback Dolphin), March 2025.
- Conservation Advice for Glyphis glyphis (Speartooth Shark), April 2014.
- Conservation Advice for Glyphis garricki (Northern River Shark). April 2014.
- Conservation Advice for Pristis pristis (Largetooth Sawfish), April 2014.
- Conservation Advice for Pristis zijsron (Green Sawfish), undated.
- Conservation Advice for Pristis clavata (Dwarf Sawfish), October 2009.

Species Recovery Plans:

- Sawfish & River Sharks Multi-species Recovery Plan, 2015.
- Recovery Plan for Marine Turtles in Australia, 2017-2027.

Threat Abatement Plans:

• Threat Abatement Plan for the impacts of marine debris on the vertebrate wildlife of Australia's coasts and oceans (2018).

ANNEX 1: DCCEEW RFI

NOTE: Attachment B: Comments on the Environmental Management Plan (EMP) - Draft 1 is addressed separately in EPBC Supplementary Report No. 3 - Environmental Management Plan Draft 2.